

# Stochastic regularization effects of semi-martingales on random functions

Romain Duboscq<sup>1</sup>   Anthony Réveillac<sup>1</sup>

<sup>1</sup>INSA de Toulouse, Institut de Mathématiques de Toulouse

London Paris Bachelier Workshop,

London, 25-26 September 2015

Stochastic regularization

Itô-Wentzell-Tanaka trick

## Stochastic regularization in a nutshell

The following slides are based on the lecture notes of Franco Flandoli (2015) and on his St. Flour lecture Notes "Random Perturbation of PDEs and Fluid Dynamic Models" (2010).

## A first example

- Let  $\varphi : \mathbb{R}^d \rightarrow \mathbb{R}$  be a **non smooth** bounded measurable map

## A first example

- Let  $\varphi : \mathbb{R}^d \rightarrow \mathbb{R}$  be a **non smooth** bounded measurable map
- Perturb it by adding a Brownian motion  $(B_t)_t$  as:

$$\varphi(x + B_t)$$

## A first example

- Let  $\varphi : \mathbb{R}^d \rightarrow \mathbb{R}$  be a **non smooth** bounded measurable map
- Perturb it by adding a Brownian motion  $(B_t)_t$  as:

$$\varphi(x + B_t)$$

- Take expectation and set:

$$u(t, x) := \mathbb{E}[\varphi(x + B_t)]$$

## A first example

- Let  $\varphi : \mathbb{R}^d \rightarrow \mathbb{R}$  be a **non smooth** bounded measurable map
- Perturb it by adding a Brownian motion  $(B_t)_t$  as:

$$\varphi(x + B_t)$$

- Take expectation and set:

$$u(t, x) := \mathbb{E}[\varphi(x + B_t)]$$

The map  $u$  is smooth and solves the Heat equation:

$$\frac{\partial u}{\partial t} = \frac{1}{2} \Delta u, \quad u(0, \cdot) = \varphi(\cdot),$$

and

$$u(t, x) = \int_{\mathbb{R}^d} P_t^{\text{heat}}(x - y) \varphi(y) dy.$$

## A second example

- Consider the following ODE:

$$dX_t = b(t, X_t)dt, \quad X_0 = x_0,$$

for some  $b : [0, T] \times \mathbb{R}^d \rightarrow \mathbb{R}^d$ .



## A second example

- Consider the following ODE:

$$dX_t = b(t, X_t)dt, \quad X_0 = x_0,$$

for some  $b : [0, T] \times \mathbb{R}^d \rightarrow \mathbb{R}^d$ .

- When  $b$  is not smooth, uniqueness may fail...

## A second example

- Consider the following ODE:

$$dX_t = b(t, X_t)dt, \quad X_0 = x_0,$$

for some  $b : [0, T] \times \mathbb{R}^d \rightarrow \mathbb{R}^d$ .

- When  $b$  is not smooth, uniqueness may fail...
- Take for instance  $d = 1$  and  $b(t, x) := b(x) := 2\operatorname{sgn}(x)\sqrt{|x|}$  and  $x_0 := 0$ , then every function of the form

$$X_t := \pm(t - t_0)^2 \mathbf{1}_{t \geq t_0}, \quad t_0 \geq 0$$

is solution.

## A second example

- Consider the following ODE:

$$dX_t = b(t, X_t)dt, \quad X_0 = x_0,$$

for some  $b : [0, T] \times \mathbb{R}^d \rightarrow \mathbb{R}^d$ .

- When  $b$  is not smooth, uniqueness may fail...
- Take for instance  $d = 1$  and  $b(t, x) := b(x) := 2\operatorname{sgn}(x)\sqrt{|x|}$  and  $x_0 := 0$ , then every function of the form

$$X_t := \pm(t - t_0)^2 \mathbf{1}_{t \geq t_0}, \quad t_0 \geq 0$$

is solution.

What is then a good solution?

## A second example

- Add some noise:

$$dX_t = b(t, X_t)dt + \sigma dB_t, \quad X_0 = x_0,$$

where  $(B_t)_t$  is a Brownian motion and  $\sigma > 0$ .

## A second example

- Add some noise:

$$dX_t = b(t, X_t)dt + \sigma dB_t, \quad X_0 = x_0,$$

where  $(B_t)_t$  is a Brownian motion and  $\sigma > 0$ .

- Why is it useful?

## A second example

- Add some noise:

$$dX_t = b(t, X_t)dt + \sigma dB_t, \quad X_0 = x_0,$$

where  $(B_t)_t$  is a Brownian motion and  $\sigma > 0$ .

- Why is it useful?
- **Selection of solutions:** Assume that for any  $\sigma$  there exists a unique solution, then let  $\mathbb{P}_\sigma$  denotes its law. Then prove that  $(\mathbb{P}_\sigma)_{\sigma>0}$  is tight and converges in law (as  $\sigma$  tends to 0) to some measure supported on the set of solutions to the ODE.

For instance, Bafico and Baldi (81') proved that for  $b(x) = 2\text{sgn}(x)\sqrt{|x|}$  and  $x_0 = 0$  it converges to:

$$\frac{1}{2}\delta_{+t^2} + \frac{1}{2}\delta_{-t^2}.$$

## A second example

- Add some noise:

$$dX_t = b(t, X_t)dt + \sigma dB_t, \quad X_0 = x_0,$$

where  $(B_t)_t$  is a Brownian motion and  $\sigma > 0$ .

- (Veretennikov 81') If  $b$  is bounded then the equation admits pathwise uniqueness.

## A second example

- Add some noise:

$$dX_t = b(t, X_t)dt + \sigma dB_t, \quad X_0 = x_0,$$

where  $(B_t)_t$  is a Brownian motion and  $\sigma > 0$ .

- (Veretennikov 81') If  $b$  is bounded then the equation admits pathwise uniqueness.
- (Krylov-Röckner 05') If  $b$  belongs to  $L^q([0, T]; L^p(\mathbb{R}^d))$  with  $\frac{d}{p} + \frac{2}{q} < 1$  ( $p, q \geq 2$ ) then the equation admits pathwise uniqueness.



## A second example

- Add some noise:

$$dX_t = b(t, X_t)dt + \sigma dB_t, \quad X_0 = x_0,$$

where  $(B_t)_t$  is a Brownian motion and  $\sigma > 0$ .

- (Veretennikov 81') If  $b$  is bounded then the equation admits pathwise uniqueness.
- (Krylov-Röckner 05') If  $b$  belongs to  $L^q([0, T]; L^p(\mathbb{R}^d))$  with  $\frac{d}{p} + \frac{2}{q} < 1$  ( $p, q \geq 2$ ) then the equation admits pathwise uniqueness.
- How does it work?

## A second example

- Recall that:

$$X_t = x_0 + \int_0^t b(s, X_s) ds + \sigma B_t$$

We try to get regularity of the blue term using the Itô-Tanaka Trick.

## A second example

- Recall that:

$$X_t = x_0 + \int_0^t b(s, X_s) ds + \sigma B_t$$

We try to get regularity of the blue term using the Itô-Tanaka Trick.

- Example: use the celebrated Itô-Tanaka formula for  $b = \delta_a$  and for  $B$ :

$$\int_0^t \delta_a(B_s) ds = |B_t - a| - |a| - \int_0^t \operatorname{sgn}(B_s - a) dB_s.$$

## A second example

- Recall that:

$$X_t = x_0 + \int_0^t b(s, X_s) ds + \sigma B_t$$

We try to get regularity of the blue term using the Itô-Tanaka Trick.

- Example: use the celebrated Itô-Tanaka formula for  $b = \delta_a$  and for  $B$ :

$$\int_0^t \delta_a(B_s) ds = |B_t - a| - |a| - \int_0^t \operatorname{sgn}(B_s - a) dB_s.$$

- Idea: to express  $\int_0^t b(s, X_s) ds$  by means of more regular objects

## The Itô-Tanaka trick

- Apply Itô's formula with a smooth mapping  $U$ :

$$U(t, X_t) = U(T, X_T) - \int_t^T \left( \frac{\partial U}{\partial t} + b \cdot \nabla U + \frac{1}{2} \sigma^2 \Delta U \right) (s, X_s) ds \\ - \sigma \int_t^T \nabla U(s, X_s) dB_s$$

## The Itô-Tanaka trick

- Apply Itô's formula with a smooth mapping  $U$ :

$$U(t, X_t) = U(T, X_T) - \int_t^T \left( \frac{\partial U}{\partial t} + b \cdot \nabla U + \frac{1}{2} \sigma^2 \Delta U \right) (s, X_s) ds \\ - \sigma \int_t^T \nabla U(s, X_s) dB_s$$

- So if  $U$  is solution to the Fokker-Planck (Backward) PDE

$$\frac{\partial U}{\partial t} + b \cdot \nabla U + \frac{\sigma^2}{2} \Delta U = -b, \quad U(T, x) = 0,$$

then

$$\int_t^T b(s, X_s) ds = -U(t, X_t) + \sigma \int_t^T \nabla U(s, X_s) dB_s$$

and so

$$X_t = x_0 + U(0, x_0) - U(t, X_t) + \sigma \int_0^t (\nabla U(s, X_s) + Id.) dB_s.$$

## Applications of the Itô-Tanaka trick to SPDEs

- The Itô-Tanaka Trick can be used to obtain new results in linear transport equations by introducing a stochastic perturbation (see *Flandoli, Gubinelli, Priola; 10'; Invent. Math.*).

## Applications of the Itô-Tanaka trick to SPDEs

- The Itô-Tanaka Trick can be used to obtain new results in linear transport equations by introducing a stochastic perturbation (see *Flandoli, Gubinelli, Priola; 10'; Invent. Math.*).
- Limitation to other problems: ([Flandoli et al.](#))

*"The generalization to nonlinear transport equations, where  $b$  depends on  $u$  itself, would be a major next step for applications to fluid dynamics but it turns out to be a difficult problem. Specifically there are already some difficulties in dealing with a vector field  $b$  which depends itself on the random perturbation  $W$ . There is no obvious extension of the Itô-Tanaka trick to integrals of the form  $\int_0^T f(\omega, s, X_s^x(\omega)) ds$  with random  $f$ ."*



Stochastic regularization

Itô-Wentzell-Tanaka trick

## Generalizations to random mappings

The problem pointed out previously is to provide an expression for:

$$\int_0^T f(s, \omega, X_s) ds,$$

where  $f$  is now random (previously we had  $f = b$  where  $b$  was deterministic) in a predictable way.

- If we reproduce the ideas before we need to consider the Fokker-Planck SPDE:

$$U(t, x) = - \int_t^T \left( \frac{1}{2} \Delta + b(s, \omega, x) \cdot \nabla \right) U(s, x) ds - \int_t^T f(s, \omega, x) ds.$$

## Generalizations to random mappings

The problem pointed out previously is to provide an expression for:

$$\int_0^T f(s, \omega, X_s) ds,$$

where  $f$  is now random (previously we had  $f = b$  where  $b$  was deterministic) in a predictable way.

- If we reproduce the ideas before we need to consider the Fokker-Planck SPDE:

$$U(t, x) = - \int_t^T \left( \frac{1}{2} \Delta + b(s, \omega, x) \cdot \nabla \right) U(s, x) ds - \int_t^T f(s, \omega, x) ds.$$

- **But:** in that case  $U(t, x)$  is not adapted (even if the data  $b, f$  are adapted) so you can not use classical Itô calculus and the previous approach fails.

## Generalizations to random mappings

- **Idea:** make it adapted, and consider rather the following Fokker-Planck **BSPDE**:

$$U^a(t, x) = - \int_t^T \mathcal{L}_s U^a(s, x) ds - \int_t^T f(s, \omega, x) ds - \int_t^T Z(s, x) dB_s,$$

$$\text{with } \mathcal{L}_s := \frac{1}{2} \Delta + b(s, \omega, x) \cdot \nabla.$$

If solvable,  $U^a$  and  $Z$  are two predictable processes.

# Itô-Wentzell-Tanaka trick

## Theorem (Duboscq, R.)

Assume that  $U^a$  and  $Z$  exist and are regular enough, then

$$\int_0^T f(s, \omega, X_s) ds = -U^a(0, X_0) - \int_0^T (\nabla U^a(s, X_s) + Z(s, X_s)) dB_s \\ - \int_0^T \nabla Z(s, X_s) ds, \mathbb{P} - a.s..$$

# Itô-Wentzell-Tanaka trick

## Theorem (Duboscq, R.)

*Assume that  $U^a$  and  $Z$  exist and are regular enough, then*

$$\int_0^T f(s, \omega, X_s) ds = -U^a(0, X_0) - \int_0^T (\nabla U^a(s, X_s) + Z(s, X_s)) dB_s \\ - \int_0^T \nabla Z(s, X_s) ds, \mathbb{P} - a.s..$$

Now we need to study the BSPDE and the regularity of  $(U^a, Z)$ .

## Analysis of the BSPDE

### Theorem (Duboscq, R.)

Let  $p, q \geq 2$ . Assume that  $b, f$  are adapted and that  $f$  belongs to a " $L^p - L^q$  space" and is Malliavin differentiable. There exists a unique strong (predictable) solution to the Fokker-Planck BSPDE

$$(U^a, Z) \in ("L^p - L^q \text{ space}")^2.$$

Futhermore, we have the following representation of  $U^a$

$$U^a(t, x) = \mathbb{E} \left[ - \int_t^T P_{t,r}^X f(r, x) dr \middle| \mathcal{F}_t \right]. \quad (1)$$

In addition, for a.e.  $(t, x)$ ,  $U^a(t, x)$  is Malliavin differentiable, and for a.e.  $x \in \mathbb{R}^d$ , a version of the process  $(Z(t, x))_{t \in [0, T]}$  is given by

$$Z(t, x) = D_t U^a(t, x) = \mathbb{E} \left[ - \int_t^T D_t P_{t,r}^X f(r, x) dr \middle| \mathcal{F}_t \right]. \quad (2)$$

# Analysis of the BSPDE

## Theorem (Duboscq, R.)

... Finally,  $U^a$  admits the following mild (a.k.a. Duhamel's formula) representation

$$U^a(t, x) = - \int_t^T P_{t,r}^X f(r, x) dr - \int_t^T P_{t,r}^X Z(r, x) dB_r, \quad (3)$$

where  $P_{s,t}^X \phi$  is the unique solution to:

$$P_{s,t}^X \phi(x) = \phi(x) - \int_s^t \mathcal{L}_r P_{r,t}^X \phi(x) dr, \quad 0 \leq s \leq t.$$



# Analysis of the BSPDE

## Remarks

- We are not working in  $L^2$
- We provide an explicit representation which is a counterpart of the one for linear BSDEs (**no reversibility of the semigroup**)
- Malliavin differentiability in  $L^p - L^q$  spaces is not completely trivial...there are catches
- Duhamel's formula in that context is new