

Civ.Eng. 2 Mathematics: Directional Derivatives

This sheet can be found on the Web: <http://www.ma.ic.ac.uk/~ajm8/Civ2>

A function f which associates a scalar value $f(\mathbf{r}) \equiv f(x, y, z)$ with every position vector $\mathbf{r} \equiv x\mathbf{i} + y\mathbf{j} + z\mathbf{k}$ is said to be a **SCALAR FIELD**, e.g. pressure, temperature.

$\frac{\partial f}{\partial x}(\mathbf{r})$, $\frac{\partial f}{\partial y}(\mathbf{r})$ and $\frac{\partial f}{\partial z}(\mathbf{r})$ are the rates of change of f at the point \mathbf{r} in the directions \mathbf{i} , \mathbf{j} and \mathbf{k} , respectively.

One may be interested in the rate of change of f at the point \mathbf{r} in some other direction. The rate of change of f at \mathbf{r} in the direction of the unit vector $\hat{\mathbf{a}}$ is said to be the **DIRECTIONAL DERIVATIVE** of f at \mathbf{r} in the direction $\hat{\mathbf{a}}$. This generalises the idea of a partial derivative.

CALCULATION of a DIRECTIONAL DERIVATIVE

Let the point P have position vector $\mathbf{p} \equiv p_1\mathbf{i} + p_2\mathbf{j} + p_3\mathbf{k}$ and let $\hat{\mathbf{a}} \equiv \hat{a}_1\mathbf{i} + \hat{a}_2\mathbf{j} + \hat{a}_3\mathbf{k}$. Then a point a distance s from P in the direction of $\hat{\mathbf{a}}$ has position vector

$$\mathbf{r} \equiv x(s)\mathbf{i} + y(s)\mathbf{j} + z(s)\mathbf{k} = \mathbf{p} + s\hat{\mathbf{a}} = (p_1 + s\hat{a}_1)\mathbf{i} + (p_2 + s\hat{a}_2)\mathbf{j} + (p_3 + s\hat{a}_3)\mathbf{k}$$

The derivative of f in the direction of $\hat{\mathbf{a}}$ is therefore

$$\begin{aligned} \lim_{s \rightarrow 0} \left[\frac{f(\mathbf{r}) - f(\mathbf{p})}{|\mathbf{r} - \mathbf{p}|} \right] &= \lim_{s \rightarrow 0} \left[\frac{f(x(s), y(s), z(s)) - f(x(0), y(0), z(0))}{s} \right] \\ &= \lim_{s \rightarrow 0} \left[\frac{F(s) - F(0)}{s} \right] = \frac{dF}{ds}(0), \end{aligned} \quad (1)$$

where $F(s) \equiv f(x(s), y(s), z(s))$. From the CHAIN RULE we have that

$$\begin{aligned} \frac{dF}{ds}(s) &= \frac{\partial f}{\partial x} \frac{dx}{ds} + \frac{\partial f}{\partial y} \frac{dy}{ds} + \frac{\partial f}{\partial z} \frac{dz}{ds} \\ &= \hat{a}_1 \frac{\partial f}{\partial x} + \hat{a}_2 \frac{\partial f}{\partial y} + \hat{a}_3 \frac{\partial f}{\partial z}. \end{aligned} \quad (2)$$

Introducing

$$\text{grad } f(\mathbf{r}) \equiv \nabla f(\mathbf{r}) \equiv \frac{\partial f}{\partial x}(\mathbf{r}) \mathbf{i} + \frac{\partial f}{\partial y}(\mathbf{r}) \mathbf{j} + \frac{\partial f}{\partial z}(\mathbf{r}) \mathbf{k}; \quad (3)$$

then from (1) and (2) it follows that

$$\begin{aligned} &\text{the directional derivative of } f \text{ at } \mathbf{p} \text{ in the direction } \hat{\mathbf{a}} = \hat{\mathbf{a}} \cdot \nabla f(\mathbf{p}), \\ \implies &\text{the directional derivative of } f \text{ at } \mathbf{r} \text{ in the direction } \hat{\mathbf{a}} = \hat{\mathbf{a}} \cdot \nabla f(\mathbf{r}), \end{aligned} \quad (4)$$

where \mathbf{r} now denotes a general point. Note that (4) yields that

$$\mathbf{i} \cdot \nabla f \equiv \frac{\partial f}{\partial x}, \quad \mathbf{j} \cdot \nabla f \equiv \frac{\partial f}{\partial y}, \quad \mathbf{k} \cdot \nabla f \equiv \frac{\partial f}{\partial z}. \quad (5)$$

$\nabla \equiv \mathbf{i} \frac{\partial}{\partial x} + \mathbf{j} \frac{\partial}{\partial y} + \mathbf{k} \frac{\partial}{\partial z}$ is a **VECTOR OPERATOR**, ∇ : scalar \rightarrow vector.

Now

$$\hat{\mathbf{a}} \cdot \nabla f = |\hat{\mathbf{a}}| |\nabla f| \cos \theta = |\nabla f| \cos \theta, \quad (6)$$

where θ is the angle between $\hat{\mathbf{a}}$ and ∇f at the point \mathbf{r} . So choosing $\hat{\mathbf{a}}$ such that $\theta = 0$ yields that

$$\text{the direction of most rapid change for } f \text{ at the point } \mathbf{r} = \frac{\nabla f}{|\nabla f|}, \quad (7)$$

$$\text{with rate of change} = |\nabla f|. \quad (8)$$

Example: Let $f(x, y, z) = 3x^2 + xy - z$. What is the rate of change of f in the direction $\mathbf{i} + 2\mathbf{j} + 3\mathbf{k}$ at the point $(1, 1, 4)$? In what direction is f changing most rapidly in at $(1, 1, 4)$?

(4), (7) and (8) yield that

$$\nabla f(x, y, z) = (6x + y) \mathbf{i} + x \mathbf{j} - \mathbf{k} \implies \nabla f(1, 1, 4) = 7\mathbf{i} + \mathbf{j} - \mathbf{k} \quad (9)$$

$$\text{unit vector in the direction } \mathbf{i} + 2\mathbf{j} + 3\mathbf{k} \text{ is } \hat{\mathbf{a}} = \frac{1}{\sqrt{14}}(\mathbf{i} + 2\mathbf{j} + 3\mathbf{k}) \implies$$

$$\text{the rate of change of } f \text{ in the direction of } \hat{\mathbf{a}} \text{ at } (1, 1, 4) = \hat{\mathbf{a}} \cdot \nabla f(1, 1, 4) = \frac{6}{\sqrt{14}} \approx 1.6,$$

$$\text{the direction of most rapid change for } f \text{ at } (1, 1, 4) = \frac{\nabla f(1, 1, 4)}{|\nabla f(1, 1, 4)|} = \frac{1}{\sqrt{51}}(7\mathbf{i} + \mathbf{j} - \mathbf{k})$$

$$\text{with rate of change} = |\nabla f(1, 1, 4)| = \sqrt{51} \approx 7.14. \quad (10)$$

The curves $f(x, y) = \text{constant}$ are called level curves (contours) of f . The surfaces $f(x, y, z) = \text{constant}$ are called level surfaces of f , e.g. isotherms, isobars, equipotentials etc.

Let C be a level curve of $f(x, y)$, parameterised by $\mathbf{r}(t) = x(t)\mathbf{i} + y(t)\mathbf{j}$; that is,

$$F(t) \equiv f(x(t), y(t)) = \text{constant} \implies 0 = \frac{dF}{dt} = \frac{\partial f}{\partial x} \frac{dx}{dt} + \frac{\partial f}{\partial y} \frac{dy}{dt} \implies \nabla f(\mathbf{r}(t)) \cdot \frac{d\mathbf{r}}{dt}. \quad (11)$$

Recall $\frac{d\mathbf{r}}{dt}(t) = \frac{dx}{dt}(t)\mathbf{i} + \frac{dy}{dt}(t)\mathbf{j}$ is tangential to C at $\mathbf{r}(t)$. Therefore (11) yields that $\nabla f(\mathbf{r}(t))$ is perpendicular (normal) to C at $\mathbf{r}(t)$. Similarly, in three space dimensions $\nabla f(\mathbf{r})$ is perpendicular (normal) to the level surface passing through \mathbf{r} .

Example: Let $f(x, y, z) = 3x^2 + xy - z$. Find the equation of the tangent plane to the level surface $f(x, y, z) = 0$ at $(1, 1, 4)$.

The above and (9) yield that $\mathbf{n} = \nabla f(1, 1, 4) = 7\mathbf{i} + \mathbf{j} - \mathbf{k}$ is the normal to the tangent plane. As $\mathbf{p} = \mathbf{i} + \mathbf{j} + 4\mathbf{k}$ is the position vector of the known point on the plane, a general point on the plane, $\mathbf{r} = x\mathbf{i} + y\mathbf{j} + z\mathbf{k}$, is such that $(\mathbf{r} - \mathbf{p}) \cdot \mathbf{n} = 0 \implies 7x + y - z = 4$.

In practice one often meets scalar fields that depend only on $r = |\mathbf{r}| = \sqrt{x^2 + y^2 + z^2}$, the distance from the origin. Then

$$\nabla f(r) = \frac{df}{dr}(r) \left(\frac{\partial r}{\partial x} \mathbf{i} + \frac{\partial r}{\partial y} \mathbf{j} + \frac{\partial r}{\partial z} \mathbf{k} \right) = \frac{1}{r} \frac{df}{dr}(r) (x\mathbf{i} + y\mathbf{j} + z\mathbf{k}) = \frac{1}{r} \frac{df}{dr}(r) \mathbf{r}, \quad (12)$$

since

$$\frac{\partial}{\partial x}(r^2) = \frac{\partial}{\partial x}(x^2 + y^2 + z^2) \implies 2r \frac{\partial r}{\partial x} = 2x \implies \frac{\partial r}{\partial x} = \frac{x}{r} \quad \text{etc.}$$

For example if $f(r) \equiv \ln r$, then (12) yields that $\nabla f(r) = \frac{1}{r^2} \mathbf{r}$.