

1. Consider the function

$$f(x) = \frac{x^3 - 1}{x^3 + 1}.$$

- (i) Put  $f(x)$  in partial fraction form;
- (ii) Find and classify all the stationary points of  $f(x)$ ;
- (iii) Find all the points of inflexion;
- (iv) Sketch the graph of  $f(x)$ , carefully indicating all the important features on your sketch (including the stationary points and points of inflexion);
- (v) Write  $f(x)$  as the sum of an odd function of  $x$  and an even function of  $x$ .

2. If

$$\zeta = \cot\left(\frac{\theta}{2}\right) e^{i\phi}$$

where  $\theta$  and  $\phi$  are real, show that

$$\begin{aligned}\sin \theta &= \frac{2\sqrt{\zeta\bar{\zeta}}}{1 + \zeta\bar{\zeta}}, \\ \cos \theta &= \frac{\zeta\bar{\zeta} - 1}{\zeta\bar{\zeta} + 1}.\end{aligned}$$

Let  $z$  be some complex number. Show that the transformation

$$z \mapsto \frac{z + 2}{z - 2}$$

maps the unit circle in the  $z$ -plane to another circle and find the centre and radius of this circle.

Sketch the locus of the curve given by  $|z - 1| = 2|z + 1|$ .

3. Find the following integrals:

(a)

$$\int \frac{dx}{x^3 - x^2 + 2x - 2};$$

(b)

$$\int \frac{dx}{e^{2x} + 1};$$

(c)

$$\int \frac{dx}{\cos x - \sqrt{2} \sin x};$$

(d)

$$\int \tanh^{-1} x \, dx.$$

Let

$$I_n = \int_0^{\pi/4} \sec^n x \, dx.$$

Determine a recurrence relation relating  $I_n$  and  $I_{n-2}$  for  $n \geq 2$ .

4. Consider the function  $\tanh^{-1} x$ .

- (a) Find an expression for  $\tanh^{-1} x$  in terms of logarithms;
- (b) Find an expression for the  $n$ -th derivative of  $\tanh^{-1} x$  where  $n \geq 1$  is any integer;
- (c) Find the complete Taylor expansion of  $\tanh^{-1} x$  about  $x = 0$ ;
- (d) Suppose the approximation

$$\tanh^{-1} x \approx x$$

is used to approximate the function  $\tanh^{-1} x$  in the interval  $0 \leq x \leq 1/2$ , use Taylor's theorem to find an estimate of the maximum possible error incurred in using this approximation.

5. Find the general solutions of the following ordinary differential equations:

(a)

$$\frac{dy}{dt} - \frac{ty}{t^2 + 1} = 1;$$

(b)

$$\frac{dy}{dt} = -1 - \frac{2t}{y - t}.$$

Find the solution of the coupled system of ordinary differential equations

$$\begin{aligned}\frac{dx}{dt} + yx &= 1, \\ \frac{dy}{dt} + \frac{y}{t} &= 0\end{aligned}$$

with  $x(1) = 1, y(1) = 2$ .