

## M2PM2 Algebra II: Solutions to Problem Sheet 8

1. Define  $A \sim B$  if  $\exists P$  invertible such that  $B = P^{-1}AP$ .

Then  $A \sim A$  as  $A = I^{-1}AI$ .

And  $A \sim B \Rightarrow B = P^{-1}AP \Rightarrow A = PBP^{-1} \Rightarrow B \sim A$ .

Finally  $A \sim B, B \sim C \Rightarrow B = P^{-1}AP, C = Q^{-1}BQ \Rightarrow C = Q^{-1}P^{-1}APQ = (PQ)^{-1}A(PQ) \Rightarrow A \sim C$ .

Hence  $\sim$  is an equivalence relation.

2. Routine first year stuff:  $P = \begin{pmatrix} 1 & 2 \\ 3 & 5 \end{pmatrix}$ ,  $Q = \begin{pmatrix} -5 & 2 \\ 3 & -1 \end{pmatrix}$ ,  $[v]_E = (a, b)^T$ ,  $[v]_F = (-5a + 2b, 3a - b)^T$ ,  $[T]_E = \begin{pmatrix} 0 & 2 \\ 3 & -1 \end{pmatrix}$ ,  $[T]_F = \begin{pmatrix} -30 & -48 \\ 18 & 29 \end{pmatrix}$ .

3. (i) The determinant must be 0 because  $T$  is not surjective (one spots  $(x_1 - x_2 + 2x_3) + (-x_1 - 3x_3) + (x_2 + x_3) = 0$ ) and hence, by rank-nullity, can't be injective either. Alternatively, just bash it out.

- (ii) The matrix of  $T$  w.r.t the usual basis  $1, x, x^2, x^3$  is triangular with diagonal entries all 1, so has determinant 1.

- (iii) Matrix of  $T$  w.r.t. basis  $\begin{pmatrix} 1 & 0 \\ 0 & 0 \end{pmatrix}, \begin{pmatrix} 0 & 0 \\ 1 & 0 \end{pmatrix}, \begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix}, \begin{pmatrix} 0 & 0 \\ 0 & 1 \end{pmatrix}$  is  $A = \begin{pmatrix} 1 & -2 & 0 & 0 \\ 1 & 4 & 0 & 0 \\ 0 & 0 & 1 & -2 \\ 0 & 0 & 1 & 4 \end{pmatrix}$ , which has determinant equal to  $(\det(M))^2 = 36$ .

4. The  $T$  of Q3(ii) satisfies  $T(1) = 1$ ,  $T(x) = x$ ,  $T(x^2) = x^2 + 4x - 1$ , and  $T(x^3) = x^3 + 9x - 2$ , so matrix of  $T$  w.r.t. basis  $1, x, x^2, x^3$  is

$$\begin{pmatrix} 1 & 0 & -1 & -2 \\ 0 & 1 & 4 & 9 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix}$$

The only eigenvalue is 1, and basis for 1-eigenspace of  $T$  is checked to be  $1, x$  without too much trouble. There is hence no basis of eigenvectors:  $g(1) < a(1)$ .

The  $T$  of Q3(iii) has matrix  $A$  as in the solution above. The characteristic polynomial of the matrix  $\begin{pmatrix} 1 & -2 \\ 1 & 4 \end{pmatrix}$  is  $(x - 2)(x - 3)$  so this matrix has distinct eigenvalues so can be diagonalised, say by a  $2 \times 2$  matrix  $P$ . Then the  $4 \times 4$  matrix  $\begin{pmatrix} P & 0 \\ 0 & P \end{pmatrix}$  diagonalises  $A$ . The eigenspaces are  $\begin{pmatrix} -2a & -2b \\ a & b \end{pmatrix}$  and  $\begin{pmatrix} a & b \\ -a & -b \end{pmatrix}$ .

5. (a)(i) Characteristic polynomial is  $(x + 1)^2(x - 2)$ , so eigenvalues are  $-1, 2$  with algebraic multiplicities 2,1 respectively. The geometric multiplicity of the eigenvalue  $-1$  is dimension of the  $-1$  eigenspace, which is easily checked to be 1; the geometric

multiplicity of 2 must also be 1 (as  $1 \leq g(2) \leq 1$ ). Since the geom multiplicity of  $-1$  is less than the algebraic multiplicity, there is no basis of eigenvectors.

(ii)  $T$  sends  $1 \rightarrow 0$ ,  $x \rightarrow 3x$ ,  $x^2 \rightarrow x + 6x^2$ , so matrix of  $T$  wrt basis  $1, x, x^2$  is  $\begin{pmatrix} 0 & 0 & 0 \\ 0 & 3 & 1 \\ 0 & 0 & 6 \end{pmatrix}$ . This has distinct eigenvalues 0,3,6, all with algebraic and geometric multiplicity 1, and there is a basis of eigenvectors.

(b) The char poly is  $(x + 1)^2(x - 1)$ , so (as in part (a)(i) above)  $A$  is diagonalisable iff the  $-1$  eigenspace has dimension 2. This eigenspace consists of solutions to the system  $\begin{pmatrix} 0 & a & b \\ 0 & 2 & c \\ 0 & 0 & 0 \end{pmatrix} v = 0$ , so it is 2-dimensional iff  $ac - 2b = 0$ .

**6.** (i) Well-definedness of multiplication (i.e. “closure”):  $S, T \in GL(V)$  implies that  $ST$  is a linear transformation, and it is invertible as  $(ST)^{-1} = T^{-1}S^{-1}$ , so  $ST \in GL(V)$ .

Associativity: follows from associativity of composition.

Identity: is identity map  $I(v) = v \forall v \in V$ .

Inverse: exists by defition.

Hence  $GL(V)$  is a group.

(ii) If  $T, U \in GL(V)$  then  $\det(TU) = \det(T)\det(U)$  by lectures, so  $\det$  is a homomorphism. Let  $B = \{v_1, \dots, v_n\}$  be a basis of  $V$ . For  $\lambda \in \mathbb{R}^*$ , define  $T : V \rightarrow V$  to be the linear map which sends

$$v_1 \rightarrow \lambda v_1, v_2 \rightarrow v_2, \dots, v_n \rightarrow v_n.$$

Then  $\det(T) = \lambda$ . Hence  $\det$  is surjective.

(iii) Fix a basis  $B$  of  $V$ . Then the map  $T \rightarrow [T]_B$  is an isomorphism from  $GL(V)$  to  $GL(n, \mathbb{R})$ .