

EXERCISES 4

Multivariate Transformations, Limits of Moment Generating Functions, Order Statistics

1* If X and Y are continuous random variables with joint pdf $f_{X,Y}$, with

$$f_{X,Y}(x,y) = c(\alpha,\beta)x^{\alpha-1}(1-x)^{\beta-1}c(\alpha+\beta,\gamma)y^{\alpha+\beta-1}(1-y)^{\gamma-1} \quad 0 < x < 1, 0 < y < 1.$$

and zero otherwise, for $\alpha, \beta, \gamma > 0$, where normalizing constants $c(\alpha, \beta)$ and $c(\alpha + \beta, \gamma)$ are defined by

$$\frac{1}{c(a,b)} = \int_0^1 t^{a-1}(1-t)^{b-1} dt$$

Let U be defined by $U = XY$. Recalling from lecture notes that

$$f_U(u) = \int_{-\infty}^{\infty} f_{X,Y}(x, u/x) |x|^{-1} dx$$

find the marginal pdf of random variable $U = XY$. Note that, as $\mathbb{X} = \mathbb{Y} = (0, 1)$, we have $XY < X$.

2. Suppose that X and Y have joint pdf given by

$$f_{X,Y}(x,y) = cxy(1-x-y) \quad 0 < x < 1, 0 < y < 1, 0 < x+y < 1.$$

for some constant $c > 0$. Find the covariance of X and Y .

3. Suppose that X and Y have joint pdf that is *constant* on the range $\mathbb{X}^{(2)} = (0, 1) \times (0, 1)$, and zero otherwise. Find the marginal pdf of random variables $U = X/Y$ and $V = -\log(XY)$. Stating clearly the range of the transformed random variable in each case. Find the pdf and cdf of $Z = X - Y$.

4. Suppose that continuous random variables X_1, X_2, X_3 are independent, and have marginal pdfs specified by

$$f_{X_i}(x_i) = c_i x_i^i e^{-x_i} \quad x_i > 0$$

for $i = 1, 2, 3$. Find the joint pdf of random variables Y_1, Y_2, Y_3 defined by

$$Y_1 = X_1/(X_1 + X_2 + X_3) \quad Y_2 = X_2/(X_1 + X_2 + X_3) \quad Y_3 = X_3/(X_1 + X_2 + X_3)$$

and evaluate the marginal expectation of Y_1 .

5. Suppose that X and Y are continuous random variables with pdf given by

$$f_{X,Y}(x,y) = \frac{1}{2\pi} \exp\left\{-\frac{1}{2}(x^2 + y^2)\right\} \quad x, y \in \mathbb{R}$$

Let random variable U be defined by $U = X/Y$. Find the pdf of U .

Suppose now that S is a random variable, independent of X and Y , with pdf given by

$$f_S(s) = c(\nu) s^{\nu/2-1} e^{-s/2} \quad s > 0$$

where ν is a positive integer and $c(\nu)$ is a constant function of ν . Find the pdf of random variable T defined by

$$T = \frac{X}{\sqrt{S/\nu}}$$

6. Suppose that the joint pdf of random variables X and Y is specified via the conditional density $f_{X|Y}$ and the marginal density f_Y as

$$f_{X|Y}(x|y) = \sqrt{\frac{y}{2\pi}} \exp\left\{-\frac{yx^2}{2}\right\} \quad x \in \mathbb{R} \quad f_Y(y) = c(\nu)y^{\nu/2-1}e^{-\nu y/2} \quad y > 0$$

where ν is a positive integer. Find the marginal pdf of X .

7. Suppose that discrete random variable X has mass function given by

$$f_X(x) = \frac{\lambda^x e^{-\lambda}}{x!} \quad x = 0, 1, 2, \dots$$

and zero otherwise, for $\lambda > 0$, and let Z_1 be defined by $Z_1 = (X - \lambda)/\sqrt{\lambda}$. Find the mgf of Z_1 . and limiting mgf as $\lambda \rightarrow \infty$.

Now suppose that X is continuous, with pdf given by $f_X(x) = cx^{\lambda-1}e^{-x}$ for $x > 0$, and zero otherwise, for $\lambda > 0$. For Z_2 defined by $Z_2 = (X - \lambda)/\sqrt{\lambda}$, find the mgf of Z_2 , and the limiting mgf as $\lambda \rightarrow \infty$.

8. Suppose that X_1 and X_2 are independent continuous random variables with pdf f_X and cdf F_X . Let Y_1 and Y_2 be the smaller and larger of the pair $\{X_1, X_2\}$ respectively. If constant c is defined by $F_X(c) = p$, for some p where $0 \leq p \leq 1$, evaluate $P[Y_2 > c]$.

9. Suppose that X_1, \dots, X_k are i.i.d continuous random variables with density function

$$f_X(x) = \lambda e^{-\lambda x} \quad x > 0$$

and zero otherwise, for some $\lambda > 0$. Find the marginal pdf of the minimum order statistic, Y_1 .

Now suppose that X_1, \dots, X_k are i.i.d continuous random variables with density function

$$f_X(x) = 1/x^2 \quad 1 \leq x < \infty$$

Find the joint pdf of the order statistics Y_1, \dots, Y_k and the marginal pdf of the smallest order and largest order statistics Y_1 and Y_k .

10. Suppose that X_1, \dots, X_k are i.i.d. continuous random variables with pdf that is constant on $(0, 1)$, zero otherwise, and let Y_k be the corresponding largest order statistic. Find the smallest k such that $P[Y_k \geq 0.99] \geq 0.95$.

11. Suppose that X is a continuous random variable with pdf given by

$$f_X(x) = 5x^4 \quad 0 \leq x \leq 1$$

and zero otherwise. Suppose that f_X above is the pdf for the marks (out of 100, and rescaled to 0-1) for a given student in a test. If test results for k tests are regarded as i.i.d. random variables having pdf f_X , let Y_1 be the random variable corresponding to the minimum score obtained in the k tests, evaluate $P[Y_1 \leq 0.75]$ if $k = 3, 4, 5$. Find the pdf of Y_1 , and the limiting probability distribution of Y_1 as the number of tests, k becomes large.

12. Suppose that X_1, \dots, X_n are iid from pdf given by

$$f_X(x) = \frac{(\alpha + 1)x^\alpha}{\theta^{\alpha+1}} \quad 0 \leq x \leq \theta$$

and zero otherwise, for parameters $\alpha, \theta > 0$. Let Y_1, \dots, Y_n be the corresponding order statistics derived from X_1, \dots, X_n . By considering cdfs, find the distributions of Y_1 and Y_n as $n \rightarrow \infty$.