

TOTAL PROBABILITY AND BAYES THEOREM

WORKED EXAMPLES

EXAMPLE 1. A biased coin (with probability of obtaining a Head equal to $p > 0$) is tossed repeatedly and independently until the first head is observed. Compute the probability that the first head appears at an **even** numbered toss.

SOLUTION: Define

- sample space Ω to be all possible infinite binary sequences of coin tosses
- event H_1 - head on **first** toss
- event E - first head on even numbered toss.

We want $P(E)$: using the Theorem of Total Probability, and the partition of Ω given by $\{H_1, H_1'\}$

$$P(E) = P(E|H_1)P(H_1) + P(E|H_1')P(H_1')$$

Now clearly, $P(E|H_1) = 0$ (given H_1 , that a head appears on the first toss, E cannot occur) and also $P(E|H_1')$ can be seen to be given by

$$P(E|H_1') = P(E') = 1 - P(E)$$

(given that a head does **not** appear on the first toss, the required conditional probability is merely the probability that the sequence concludes after a further **odd** number of tosses, that is, the probability of E'). Hence $P(E)$ satisfies

$$P(E) = 0 \times p + (1 - P(E)) \times (1 - p) = (1 - p)(1 - P(E))$$

so that

$$P(E) = \frac{1 - p}{2 - p}.$$

Alternately, consider the partition of E into E_1, E_2, \dots where E_k is the event that the first head occurs on the $2k$ th toss. Then $E = \bigcup_{k=1}^{\infty} E_k$, and

$$P(E) = P\left(\bigcup_{k=1}^{\infty} E_k\right) = \sum_{k=1}^{\infty} P(E_k).$$

Now $P(E_k) = (1 - p)^{2k-1} p$ (that is, $2k - 1$ tails, then a head), so

$$\begin{aligned} P(E) &= \sum_{k=1}^{\infty} (1 - p)^{2k-1} p \\ &= \frac{p}{1 - p} \sum_{k=1}^{\infty} (1 - p)^{2k} \\ &= \frac{p}{1 - p} \frac{(1 - p)^2}{1 - (1 - p)^2} \\ &= \frac{1 - p}{2 - p} \end{aligned}$$

EXAMPLE 2. Two players A and B are competing at a trivia quiz game involving a series of questions. On any individual question, the probabilities that A and B give the correct answer are α and β respectively, for all questions, with outcomes for different questions being independent. The game finishes when a player wins by answering a question correctly.

Compute the probability that A wins if

- (a) A answers the first question .(b) B answers the first question

SOLUTION: Define

- sample space Ω to be all possible infinite sequences of answers
- event A - A answers the first question
- event F - game ends after the first question
- event W - A wins.

We want

$$P(W|A) \quad \text{and} \quad P(W|A')$$

Using the Theorem of Total Probability, and the partition given by $\{F, F'\}$

$$P(W|A) = P(W|A \cap F)P(F|A) + P(W|A \cap F')P(F'|A).$$

Now, clearly

$$P(F|A) = P[\text{A answers first question correctly}] = \alpha \quad P(F'|A) = 1 - \alpha$$

and $P(W|A \cap F) = 1$, but $P(W|A \cap F') = P(W|A')$, so that

$$P(W|A) = (1 \times \alpha) + (P(W|A') \times (1 - \alpha)) . = \alpha + P(W|A')(1 - \alpha) \quad (1)$$

Similarly

$$P(W|A') = P(W|A' \cap F)P(F|A') + P(W|A' \cap F')P(F'|A').$$

We have

$$P(F|A') = P[\text{B answers first question correctly}] = \beta \quad P(F'|A) = 1 - \beta$$

but $P(W|A' \cap F) = 0$. Finally $P(W|A' \cap F') = P(W|A)$, so that

$$P(W|A') = (0 \times \beta) + (P(W|A) \times (1 - \beta)) . = P(W|A)(1 - \beta) \quad (2)$$

Solving (1) and (2) simultaneously gives, for (a) and (b)

$$P(W|A) = \frac{\alpha}{1 - (1 - \alpha)(1 - \beta)} \quad P(W|A') = \frac{(1 - \beta)\alpha}{1 - (1 - \alpha)(1 - \beta)}$$

Note: recall, for any events E_1 and E_2 we have that

$$P(E'_1|E_2) = 1 - P(E_1|E_2)$$

but not necessarily that

$$P(E_1|E'_2) = 1 - P(E_1|E_2)$$

EXAMPLE 3. Patients are recruited onto the two arms (0 - Control, 1-Treatment) of a clinical trial. The probability that an adverse outcome occurs on the control arm is p_0 , and is p_1 for the treatment arm. Patients are allocated alternately onto the two arms in the sequence 010101..., and their outcomes are independent

What is the probability that the first adverse event occurs on the control arm. ?

SOLUTION: Define

- sample space Ω to be all possible infinite sequences of patients outcomes
- event E_1 - first patient (allocated onto the control arm) suffers an adverse outcome
- event E_2 - first patient (allocated onto the control arm) does not suffer an adverse outcome, but the second patient (allocated onto the treatment arm) does suffer an adverse outcome
- event E_0 - neither of the first two patients suffer adverse outcomes
- event F - first adverse event occurs on the control arm

We want $P(F)$. Now the events E_1, E_2 and E_0 partition Ω , so, by the Theorem,

$$P(F) = P(F|E_1)P(E_1) + P(F|E_2)P(E_2) + P(F|E_0)P(E_0).$$

Now

$$P(E_1) = p_0 \quad P(E_2) = (1 - p_0)p_1 \quad P(E_0) = (1 - p_0)(1 - p_1)$$

and $P(F|E_1) = 1$, $P(F|E_2) = 0$. Finally, as after two non-adverse outcomes, the allocation process effectively re-starts, so $P(F|E_0) = P(F)$. Hence

$$P(F) = (1 \times p_0) + (0 \times (1 - p_0)p_1) + (P(F) \times (1 - p_0)(1 - p_1)) = p_0 + (1 - p_0)(1 - p_1)P(F)$$

which can be re-arranged to give

$$P(F) = \frac{p_0}{p_0 + p_1 - p_0p_1}$$

EXAMPLE 4. In a tennis match, with the score at deuce, the game is one by the first player who gets a clear lead of two points.

If the probability that given player wins a particular point is θ , and all points are played independently, what is the probability that player eventually wins the game

SOLUTION: Define

- sample space Ω to be all possible infinite sequences of points
- event W_i - nominated player wins the i th point
- event V_i - nominated player wins the game on the i th point
- event V - nominated player wins the game.

We want $P(V)$. The events $\{W_1, W_1'\}$ partition Ω , and thus, by the Theorem

$$P(V) = P(V|W_1)P(W_1) + P(V|W_1')P(W_1'). \quad (3)$$

Now $P(W_1) = \theta$ and $P(W_1') = 1 - \theta$. To get $P(V|W_1)$ and $P(V|W_1')$, we need to further condition on the result of the second point, and again use the Theorem: for example

$$\begin{aligned} P(V|W_1) &= P(V|W_1 \cap W_2)P(W_2|W_1) + P(V|W_1 \cap W_2')P(W_2'|W_1) \\ P(V|W_1') &= P(V|W_1' \cap W_2)P(W_2|W_1') + P(V|W_1' \cap W_2')P(W_2'|W_1') \end{aligned} \quad (4)$$

where

$$\begin{aligned} P(V|W_1 \cap W_2) &= 1 & P(W_2|W_1) &= P(W_2) = \theta \\ P(V|W_1 \cap W_2') &= P(V) & P(W_2'|W_1) &= P(W_2') = 1 - \theta \\ P(V|W_1' \cap W_2) &= P(V) & P(W_2|W_1') &= P(W_2) = \theta \\ P(V|W_1' \cap W_2') &= 0 & P(W_2'|W_1') &= P(W_2') = 1 - \theta \end{aligned}$$

as,

given $W_1 \cap W_2$: the game is over, and the player has won
 given $W_1 \cap W_2'$: the game is back at deuce
 given $W_1' \cap W_2$: the game is back at deuce
 given $W_1' \cap W_2'$: the game is over, and the player has lost

and the results of successive points are independent. Thus

$$\begin{aligned} P(V|W_1) &= (1 \times \theta) + (P(V) \times (1 - \theta)) = \theta + (1 - \theta)P(V) \\ P(V|W_1') &= (P(V) \times \theta) + 0 \times (1 - \theta) = \theta P(V) \end{aligned}$$

Hence, combining (3) and (4) we have

$$P(V) = (\theta + (1 - \theta)P(V))\theta + \theta P(V)(1 - \theta) = \theta^2 + 2\theta(1 - \theta)P(V) \implies P(V) = \frac{\theta^2}{1 - 2\theta(1 - \theta)}$$

Alternately, $\{V_i, i = 1, 2, \dots\}$ partition V . Hence

$$P(V) = \sum_{i=1}^{\infty} P(V_i)$$

Now, $P(V_i) = 0$ if i is odd, as the game can never be completed after an odd number of points. For $i = 2$, $P(V_2) = \theta^2$, and for $i = 2k + 2$ ($k = 1, 2, 3, \dots$) we have

$$P(V_i) = P(V_{2k+2}) = 2^k \theta^k (1 - \theta)^k \times \theta^2$$

- the score must stand at deuce after $2k$ points and the game must not have been completed prior to this, indicating that there must have been k successive drawn pairs of points, each of which could be arranged win/lose or lose/win for the nominated player. Then that player must win the final two points. Hence

$$P(V) = \sum_{k=0}^{\infty} P(V_{2k+2}) = \theta^2 \sum_{k=0}^{\infty} \{2\theta(1 - \theta)\}^k = \frac{\theta^2}{1 - 2\theta(1 - \theta)}$$

as the term in the geometric series satisfies $|2\theta(1 - \theta)| < 1$.

EXAMPLE 5 A coin for which $P(\text{Heads}) = p$ is tossed until two successive Tails are obtained.

Find the probability that the experiment is completed on the n th toss.

SOLUTION: Define

- sample space Ω to be all possible infinite sequences of tosses
- event E_1 : first toss is H
- event E_2 : first two tosses are TH
- event E_3 : first two tosses are TT
- event F_n : experiment completed on the n th toss

We want $P(F_n)$ for $n = 2, 3, \dots$. The events $\{E_1, E_2, E_3\}$ partition Ω , and thus, by the Theorem

$$P(F_n) = P(F_n|E_1)P(E_1) + P(F_n|E_2)P(E_2) + P(F_n|E_3)P(E_3). \quad (5)$$

Now for $n = 2$

$$P(F_2) = P(E_3) = (1 - p)^2$$

and for $n > 2$,

$$P(F_n|E_1) = P(F_{n-1}) \quad P(F_n|E_2) = P(F_{n-2}) \quad P(F_n|E_3) = 0$$

as given E_1 we need $n - 1$ further tosses that finish TT for F_n to occur, and given E_2 , we need $n - 2$ further tosses that finish TT for F_n to occur, with all tosses independent. Hence, if $p_n = P(F_n)$ then $p_2 = (1 - p)^2$, otherwise, from (5), p_n satisfies

$$p_n = (p_{n-1} \times p) + (p_{n-2} \times (1 - p)p) = pp_{n-1} + p(1 - p)p_{n-2}$$

To find p_n explicitly, try a solution of the form $p_n = A\lambda_1^n + B\lambda_2^n$ which gives

$$A\lambda_1^n + B\lambda_2^n = p(A\lambda_1^{n-1} + B\lambda_2^{n-1}) + p(1 - p)(A\lambda_1^{n-2} + B\lambda_2^{n-2}).$$

First, collecting terms in λ_1 gives

$$\lambda_1^n = p\lambda_1^{n-1} + p(1 - p)\lambda_1^{n-2} \implies \lambda_1^2 - p\lambda_1 - p(1 - p) = 0$$

indicating that λ_1 and λ_2 are given as the roots of this quadratic, that is

$$\lambda_1 = \frac{p - \sqrt{p^2 + 4p(1 - p)}}{2} \quad \lambda_2 = \frac{p + \sqrt{p^2 + 4p(1 - p)}}{2}$$

Furthermore,

$$\begin{aligned} n = 1 : p_1 &= 0 & \implies A\lambda_1 + B\lambda_2 &= 0 \\ n = 2 : p_2 &= (1 - p)^2 & \implies A\lambda_1^2 + B\lambda_2^2 &= (1 - p)^2 \\ \implies A &= \frac{(1 - p)^2}{\lambda_1(\lambda_1 - \lambda_2)} & B &= -\frac{\lambda_1}{\lambda_2}A = -\frac{(1 - p)^2}{\lambda_2(\lambda_1 - \lambda_2)} \\ \implies p_n &= -\frac{(1 - p)^2 \lambda_1^n}{\lambda_1(\lambda_2 - \lambda_1)} + \frac{(1 - p)^2 \lambda_2^n}{\lambda_2(\lambda_2 - \lambda_1)} = \frac{(1 - p)^2}{\sqrt{p(4 - 3p)}} (\lambda_2^{n-1} - \lambda_1^{n-1}) \quad n \geq 2 \end{aligned}$$

EXAMPLE 6 Information is transmitted digitally as a binary sequence know as “bits”. However, noise on the channel corrupts the signal, in that a digit transmitted as 0 is received as 1 with probability $1 - \alpha$, with a similar random corruption when the digit 1 is transmitted. It has been observed that, across a large number of transmitted signals, the 0s and 1s are transmitted in the ratio 3 : 4.

Given that the sequence 101 is received, what is the probability distribution over transmitted signals ? Assume that the transmission and reception processes are independent

SOLUTION: Define

- sample space Ω to be all possible binary sequences of length three that is

$$\{000, 001, 010, 011, 100, 101, 110, 111\}$$

- a corresponding set of
signal events $\{S_0, S_1, S_2, S_3, S_4, S_5, S_6, S_7\}$ and
reception events $\{R_0, R_1, R_2, R_3, R_4, R_5, R_6, R_7\}$

We have observed event R_5 , that 101 was received; we wish to compute $P(S_i|R_5)$, for $i = 0, 1, \dots, 7$. However, on the information given, we only have (or can compute) $P(R_5|S_i)$. First, using the Theorem of Total Probability, we compute $P(R_5)$

$$P(R_5) = \sum_{i=0}^7 P(R_5|S_i)P(S_i). \quad (6)$$

Consider $P(R_5|S_0)$; if 000 is transmitted, the probability that 101 is received is $(1 - \alpha) \times \alpha \times (1 - \alpha) = \alpha(1 - \alpha)^2$ (corruption, no corruption, corruption) By complete evaluation we have

$$\begin{aligned} P(R_5|S_0) &= \alpha(1 - \alpha)^2 & P(R_5|S_1) &= \alpha^2(1 - \alpha) & P(R_5|S_2) &= (1 - \alpha)^3 & P(R_5|S_3) &= \alpha(1 - \alpha)^2 \\ P(R_5|S_4) &= \alpha^2(1 - \alpha) & P(R_5|S_5) &= \alpha^3 & P(R_5|S_6) &= \alpha(1 - \alpha)^2 & P(R_5|S_7) &= \alpha^2(1 - \alpha) \end{aligned}$$

Now, the prior information about digits transmitted is that the probability of transmitting a 1 is $4/7$, so

$$\begin{aligned} P(S_0) &= \left(\frac{3}{7}\right)^3 & P(S_1) &= \left(\frac{4}{7}\right)\left(\frac{3}{7}\right)^2 & P(S_2) &= \left(\frac{4}{7}\right)\left(\frac{3}{7}\right)^2 & P(S_3) &= \left(\frac{4}{7}\right)^2\left(\frac{3}{7}\right) \\ P(S_4) &= \left(\frac{4}{7}\right)\left(\frac{3}{7}\right)^2 & P(S_5) &= \left(\frac{4}{7}\right)^2\left(\frac{3}{7}\right) & P(S_6) &= \left(\frac{4}{7}\right)^2\left(\frac{3}{7}\right) & P(S_7) &= \left(\frac{4}{7}\right)^3 \end{aligned}$$

and hence (6) can be computed as

$$P(R_5) = \frac{48\alpha^3 + 136\alpha^2(1 - \alpha) + 123\alpha(1 - \alpha)^2 + 36(1 - \alpha)^3}{343}.$$

Finally, using Bayes Theorem, we have the probability distribution over $\{S_0, S_1, S_2, S_3, S_4, S_5, S_6, S_7\}$ given by

$$P(S_i|R_5) = \frac{P(R_5|S_i)P(S_i)}{P(R_5)}.$$

For example, the probability of correct reception is

$$P(S_5|R_5) = \frac{P(R_5|S_5)P(S_5)}{P(R_5)} = \frac{48\alpha^3}{48\alpha^3 + 136\alpha^2(1 - \alpha) + 123\alpha(1 - \alpha)^2 + 36(1 - \alpha)^3}$$




Figure 1: **EXAMPLE 6:** Posterior probability as a function of α

EXAMPLE 7 On a given Saturday afternoon, Oxford Street in London comprises $2/3$ Helpful people who, when asked for directions, will give the correct answer with probability $3/4$, but also $1/3$ Unhelpful people who will, maliciously or otherwise, give the incorrect answer with probability 1.

Compute the following probabilities, assuming that the answers given are conditionally independent, given the category (Helpful or Unhelpful) of person asked :

- You ask a randomly selected person whether a famous landmark is to the North or South; you are given the answer North. What is the probability that is correct ?
- You ask the same person a total of r times, and receive the same answer each time . What is the probability that it is correct ?
- If the fourth answer is, instead, South, compute the probability that North is in fact correct, in light of the first four answers
- Suppose that you believe, prior to any inquiries, that North is correct with probability p . Show that the first answer received does not contain any useful information for you.
- If the first two answers given are the same, show that still no useful information has been elicited.
- Show that, if three identical answers, the information gained should not be ignored.

SOLUTION: Define

- I_r - event that you receive r identical answers
- T - event that the answer given is correct
- H - event that the person asked is, in fact, Helpful
- N - event that the correct answer is North

We first need to compute $P(T|I_r)$ for $r = 1, 2, \dots$. Then, using the conditional probability definition, and a partition given by H ,

$$P(T|I_r) = \frac{P(T \cap I_r)}{P(I_r)} = \frac{P(T \cap I_r \cap H) + P(T \cap I_r \cap H')}{P(I_r)} = \frac{P(T \cap I_r|H)P(H)}{P(I_r)}$$

as $P(T \cap I_r \cap H') = 0$. Now

$$P(T \cap I_r|H) = \left(\frac{3}{4}\right)^r \quad P(H) = \frac{2}{3}$$

and

$$\begin{aligned} P(I_r) &= P(I_r|H)P(H) + P(I_r|H')P(H') = ([P(I_r \cap T|H) + P(I_r \cap T'|H)] \times P(H)) + (1 \times P(H')) \\ &= \left(\left[\left(\frac{3}{4}\right)^r + \left(\frac{1}{4}\right)^r \right] \times \left(\frac{2}{3}\right) \right) + \left(\frac{1}{3}\right) \end{aligned}$$

(b) Therefore, in general,

$$P(T|I_r) = \frac{\left(\frac{3}{4}\right)^r \left(\frac{2}{3}\right)}{\left[\left(\frac{3}{4}\right)^r + \left(\frac{1}{4}\right)^r \right] \left(\frac{2}{3}\right) + \left(\frac{1}{3}\right)}$$

and specifically

(a) $P(T|I_1) = \frac{1}{2}$. Note, also, that $P(T|I_2) = \frac{1}{2}$, but $P(T|I_3) = \frac{9}{20}$ and $P(T|I_4) = \frac{27}{70}$

(c) If the fourth answer is different from the previous three, then the person must be Helpful, and we can condition on H , giving

$$P(T_3 \cap D|I_3 \cap H) = \frac{P(T_3 \cap D \cap I_3|H)}{P(T_3 \cap D \cap I_3|H) + P(T_3 \cap D \cap I'_3|H)} = \frac{\left(\frac{3}{4}\right)^3 \left(\frac{1}{4}\right)}{\left(\frac{3}{4}\right)^3 \left(\frac{1}{4}\right) + \left(\frac{3}{4}\right) \left(\frac{1}{4}\right)^3} = \frac{9}{10}$$

where D means a different answer given on the fourth enquiry, T_3 means that the first 3 answers given were correct.

(d) We now want $P(\text{North Correct}|\text{Answer given is North})$ and $P(\text{North Correct}|\text{Answer given is South})$. By the conditional probability definition, utilizing a suitable partition

$$P(N_{COR}|N_{ANS}) = \frac{P(N_{COR} \cap N_{ANS})}{P(N_{ANS})}$$

Various terms that are used to construct these probabilities are

$$(1) \quad P(N_{COR} \cap N_{ANS} \cap H) = P(N_{ANS}|H \cap N_{COR})P(H|N_{COR})P(N_{COR}) = \frac{3}{4} \times \frac{2}{3} \times p$$

$$(2) \quad P(N_{COR} \cap N_{ANS} \cap H') = P(N_{ANS}|H' \cap N_{COR})P(H'|N_{COR})P(N_{COR}) = 0 \times \frac{1}{3} \times p$$

$$(3) \quad P(N'_{COR} \cap N_{ANS} \cap H) = P(N_{ANS}|H \cap N'_{COR})P(H|N'_{COR})P(N'_{COR}) = \frac{1}{4} \times \frac{2}{3} \times (1-p)$$

$$(4) \quad P(N'_{COR} \cap N_{ANS} \cap H') = P(N_{ANS}|H' \cap N'_{COR})P(H'|N'_{COR})P(N'_{COR}) = 1 \times \frac{1}{3} \times (1-p)$$

and hence

$$P(N_{COR}|N_{ANS}) = \frac{(1) + (2)}{(1) + (2) + (3) + (4)} = \frac{\frac{1}{2}p + 0}{\frac{1}{2}p + 0 + \frac{1}{6}(1-p) + \frac{1}{3}(1-p)} = p$$

It can be shown similarly that, as $N_{COR} = (N_{COR} \cap N_{ANS}) \cup (N_{COR} \cap N'_{ANS})$,

$$P(N_{COR}|N'_{ANS}) = \frac{P(N_{COR} \cap N'_{ANS})}{P(N'_{ANS})} = \frac{P(N_{COR}) - P(N_{COR} \cap N_{ANS})}{1 - P(N_{ANS})} = \frac{p - \frac{1}{2}p}{1 - \frac{1}{2}} = p$$

(e) Using an obvious extension of notation, we wish to compute

$$P(N_{COR}|N_{ANS}^{(2)}) = \frac{P(N_{COR} \cap N_{ANS}^{(2)})}{P(N_{ANS}^{(2)})}$$

for which we need

- (1) $P(N_{COR} \cap N_{ANS}^{(2)} \cap H) = P(N_{ANS}^{(2)}|H \cap N_{COR})P(H|N_{COR})P(N_{COR}) = \left(\frac{3}{4}\right)^2 \times \frac{2}{3} \times p$
- (2) $P(N_{COR} \cap N_{ANS}^{(2)} \cap H') = P(N_{ANS}^{(2)}|H' \cap N_{COR})P(H'|N_{COR})P(N_{COR}) = 0 \times \frac{1}{3} \times p$
- (3) $P(N'_{COR} \cap N_{ANS}^{(2)} \cap H) = P(N_{ANS}^{(2)}|H \cap N'_{COR})P(H|N'_{COR})P(N'_{COR}) = \left(\frac{1}{4}\right)^2 \times \frac{2}{3} \times (1-p)$
- (4) $P(N'_{COR} \cap N_{ANS}^{(2)} \cap H') = P(N_{ANS}^{(2)}|H' \cap N'_{COR})P(H'|N'_{COR})P(N'_{COR}) = 1 \times \frac{1}{3} \times (1-p)$

so that

$$P(N_{COR}|N_{ANS}^{(2)}) = \frac{(1) + (2)}{(1) + (2) + (3) + (4)} = \frac{\frac{9}{24}p + 0}{\frac{9}{24}p + 0 + \frac{1}{24}(1-p) + \frac{8}{24}(1-p)} = p$$

$$P(N_{COR}|N_{ANS}^{(2)'}) = \frac{P(N_{COR} \cap N_{ANS}^{(2)'})}{P(N_{ANS}^{(2)'})} = \frac{P(N_{COR}) - P(N_{COR} \cap N_{ANS}^{(2)})}{1 - P(N_{ANS}^{(2)})} = \frac{p - \frac{9}{24}p}{1 - \frac{9}{24}} = p$$

(f) By a further extension, first we compute

- (1) $P(N_{COR} \cap N_{ANS}^{(3)} \cap H) = P(N_{ANS}^{(3)}|H \cap N_{COR})P(H|N_{COR})P(N_{COR}) = \left(\frac{3}{4}\right)^3 \times \frac{2}{3} \times p$
- (3) $P(N_{COR} \cap N_{ANS}^{(3)} \cap H') = P(N_{ANS}^{(3)}|H' \cap N_{COR})P(H'|N_{COR})P(N_{COR}) = 0 \times \frac{1}{3} \times p$
- (3) $P(N'_{COR} \cap N_{ANS}^{(3)} \cap H) = P(N_{ANS}^{(3)}|H \cap N'_{COR})P(H|N'_{COR})P(N'_{COR}) = \left(\frac{1}{4}\right)^3 \times \frac{2}{3} \times (1-p)$
- (4) $P(N'_{COR} \cap N_{ANS}^{(3)} \cap H') = P(N_{ANS}^{(3)}|H' \cap N'_{COR})P(H'|N'_{COR})P(N'_{COR}) = 1 \times \frac{1}{3} \times (1-p)$

so that

$$P(N_{COR}|N_{ANS}^{(3)}) = \frac{(1) + (2)}{(1) + (2) + (3) + (4)} = \frac{\frac{27}{96}p + 0}{\frac{27}{96}p + 0 + \frac{1}{96}(1-p) + \frac{32}{96}(1-p)} = \frac{9p}{9p + 11(1-p)}$$

$$P(N_{COR}|N_{ANS}^{(3)'}) = \frac{P(N_{COR} \cap N_{ANS}^{(3)'})}{P(N_{ANS}^{(3)'})} = \frac{P(N_{COR}) - P(N_{COR} \cap N_{ANS}^{(3)})}{1 - P(N_{ANS}^{(3)})} = \frac{69p}{63 + 6p}$$

and the probability **has** been updated in light of both new pieces of information

EXAMPLE 8 While watching a game of Champions League football in a bar, you observe someone who is clearly supporting Manchester United in the game.

What is the probability that they were actually born within 25 miles of Manchester ?. Assume that the probability that a randomly selected person in a typical local bar environment is born within 25 miles of Manchester is $\frac{1}{20}$, and that the chance that a person born within 25 miles of Manchester actually supports United is $\frac{7}{10}$. Assume also that the probability that a person not born within 25 miles of Manchester supports United with probability $\frac{1}{10}$

SOLUTION: Define

- B - event that the person is born within 25 miles of Manchester
- U - event that the person supports United.

We want $P(B|U)$. By Bayes Theorem,

$$\begin{aligned} P(B|U) &= \frac{P(U|B)P(B)}{P(U)} = \frac{P(U|B)P(B)}{P(U|B)P(B) + P(U|B')P(B')} \\ &= \frac{\frac{7}{10} \frac{1}{20}}{\frac{7}{10} \frac{1}{20} + \frac{1}{10} \frac{19}{20}} \\ &= \frac{7}{26} \approx 0.269 \end{aligned}$$