M2S1 - EXERCISES 2

Discrete and Continuous Probability Distributions.

1. For which values of the constant c do the following functions define valid probability mass functions for a discrete random variable X, taking values on range $\mathbb{X} = \{1, 2, 3, ...\}$:

(a)
$$f_X(x) = c/2^x$$

(a)
$$f_X(x) = c/2^x$$
 (b) $f_X(x) = c/(x2^x)$

(c)
$$f_X(x) = c/(x^2)$$
 (d) $f_X(x) = c2^x/x!$

(d)
$$f_X(x) = c2^x/x$$

In each case, calculate (where possible) P[X > 1] and P[X is even]

- 2. n identical fair coins are tossed. Those that show Heads are tossed again, and the number of Heads obtained on the second set of tosses defines a discrete random variable X. Assuming that all tosses are independent, find the range and probability mass function of X.
- 3. A point is to be selected from an integer lattice restricted to the triangle with corners at (1,1), (n,1)and (n,n) for positive integer n. If all points are equally likely to be selected, find the probability mass functions for the two discrete random variables X and Y corresponding to the x- and y- coordinates of the selected points respectively.
- 4. A continuous random variable X has pdf given by

$$f_X(x) = c(1-x)x^2$$
 $0 < x < 1$

and zero otherwise. Find the value of c, the cdf of X, F_X , and P[X > 1/2].

5. A function f is defined by

$$f(x) = k/x^{k+1} \qquad x > 1$$

and zero otherwise. For what values of k is f a valid pdf? Find the cdf of X.

6. A continuous random variable X has pdf given by

$$f_X(x) = \begin{cases} x & 0 < x < 1 \\ 2 - x & 1 \le x < 2 \end{cases}$$

and zero otherwise. Sketch f_X , and find the cdf F_X .

7. A continuous random variable X has cdf given by

$$F_X(x) = c(\alpha x^{\beta} - \beta x^{\alpha}) \qquad 0 \le x \le 1$$

for constants $1 \le \beta < \alpha$, and zero otherwise. Find the value of constant c, and evaluate the rth moment of X.

A continuous random variable X has cdf given by

$$F_X(x) = \frac{2\beta x}{\beta^2 + x^2}$$
 $0 \le x \le \beta$

for constant $\beta > 0$. Find the pdf of X, and show that the expectation of X is

$$\beta(1-\log 2)$$