

**M2S1 - EXERCISES 6**  
**DISTRIBUTIONAL RESULTS**

1. The joint pdf  $f_{X,Y}$  of positive random variables  $X$  and  $Y$  is specified as

$$f_{X,Y}(x,y) = f_{X|Y}(x|y)f_Y(y)$$

where  $X|Y = y \sim \text{Exponential}(y)$  and  $Y \sim \text{Gamma}(\alpha, \beta)$ . Identify the marginal distribution of  $X$ .

2. *The Bivariate Normal Distribution:* Suppose that  $X_1$  and  $X_2$  are i.i.d  $\text{Normal}(0,1)$  random variables. Let random variables  $Y_1$  and  $Y_2$  be defined by

$$\begin{aligned} Y_1 &= \mu_1 + \sigma_1\sqrt{1-\rho^2}X_1 + \sigma_1\rho X_2 \\ Y_2 &= \mu_2 + \sigma_2 X_2 \end{aligned} \quad \text{or equivalently} \quad \begin{bmatrix} Y_1 \\ Y_2 \end{bmatrix} = \begin{bmatrix} \mu_1 \\ \mu_2 \end{bmatrix} + \begin{bmatrix} \sigma_1\sqrt{1-\rho^2} & \sigma_1\rho \\ 0 & \sigma_2 \end{bmatrix} \begin{bmatrix} X_1 \\ X_2 \end{bmatrix}$$

for positive constants  $\sigma_1$  and  $\sigma_2$ , and  $|\rho| < 1$ . Find the joint pdf of  $(Y_1, Y_2)$ .

Show that, marginally for  $i = 1, 2$ ,  $Y_i \sim \text{Normal}(\mu_i, \sigma_i^2)$ , and that conditionally

$$Y_1|Y_2 = y_2 \sim \text{Normal}\left(\mu_1 + \frac{\rho\sigma_1}{\sigma_2}(y_2 - \mu_2), \sigma_1^2(1 - \rho^2)\right)$$

$$Y_2|Y_1 = y_1 \sim \text{Normal}\left(\mu_2 + \frac{\rho\sigma_2}{\sigma_1}(y_1 - \mu_1), \sigma_2^2(1 - \rho^2)\right)$$

Find the correlation of  $Y_1$  and  $Y_2$ .

3. Suppose that  $U_1$  and  $U_2$  are i.i.d  $\text{Uniform}(0,1)$  random variables. Let random variables  $Z_1$  and  $Z_2$  be defined by

$$Z_1 = \sqrt{-2 \log U_1} \cos(2\pi U_2)$$

$$Z_2 = \sqrt{-2 \log U_1} \sin(2\pi U_2)$$

(log is the natural logarithm). Find the joint pdf of  $(Z_1, Z_2)$ .

4. Suppose that  $U$  is a  $\text{Uniform}(0,1)$  random variable. Find the distribution of

$$X = -\beta \log U.$$

Suppose that an unlimited sequence of  $\text{Uniform}(0,1)$  random variables is available. Describe how to generate

- (i) a  $\text{Gamma}(k, \lambda)$  random variable, for integer  $k \geq 0$ .
- (ii) a realization of a *Poisson process* with rate  $\mu$ .
- (iii) a *Chisquare* ( $\nu \equiv \text{Gamma}\left(\frac{\nu}{2}, \frac{1}{2}\right)$ ) random variable, where  $\nu$  is a positive, real parameter.
- (iv) a *Student*( $n$ ) random variable, where  $n$  is a positive integer parameter.

Use the results from question 3., and results given in lectures and the printed notes.