

## M2S1 : EXERCISES 2 : SOLUTIONS

1. Need  $\sum_{x=1}^{\infty} f_x(x) = 1$ . Hence

$$(a) \quad c^{-1} = \sum_{x=1}^{\infty} \frac{1}{2^x} = 1 \qquad (b) \quad c^{-1} = \sum_{x=1}^{\infty} \frac{1}{x2^x} = \log 2$$

$$(c) \quad c^{-1} = \sum_{x=1}^{\infty} \frac{1}{x^2} = \frac{\pi^2}{6} \qquad (d) \quad c^{-1} = \sum_{x=1}^{\infty} \frac{2^x}{x!} = e^2 - 1$$

(a) is given by the sum of a geometric progression; (b) uses the fact that

$$\frac{1}{1-t} = 1 + t + t^2 + \dots = \sum_{x=0}^{\infty} t^x \implies -\log(1-t) = t + \frac{t^2}{2} + \frac{t^3}{3} + \dots = \sum_{x=1}^{\infty} \frac{t^x}{x}$$

by integrating both sides with respect to  $t$ . Hence for  $t = 1/2$ , we have

$$\log 2 = -\log(1 - 1/2) = \sum_{x=1}^{\infty} \frac{1}{x2^x}.$$

(c) is a “well-known” mathematical result (**you would not be expected to remember it for the examination**); (d) uses the power series expansion of  $e^t$ , evaluated at  $t = 2$ , that is

$$e^t = \sum_{x=0}^{\infty} \frac{t^x}{x!} \implies e^2 = \sum_{x=0}^{\infty} \frac{2^x}{x!} = 1 + \sum_{x=1}^{\infty} \frac{2^x}{x!}$$

Clearly  $P[X > 1] = 1 - P[X = 1]$ , so

$$(a) \quad P[X > 1] = \frac{1}{2} \qquad (b) \quad P[X > 1] = 1 - \frac{1}{2 \log 2}$$

$$(c) \quad P[X > 1] = 1 - \frac{6}{\pi^2} \qquad (d) \quad P[X > 1] = \frac{e^2 - 3}{e^2 - 1}$$

$P[X \text{ is even}] = \sum_{x=1}^{\infty} P[X = 2i]$ , so

$$(a) \quad P[X \text{ is even}] = \frac{1}{3} \qquad (b) \quad P[X \text{ is even}] = 1 - \frac{\log 3}{\log 4}$$

$$(c) \quad P[X \text{ is even}] = \frac{1}{4} \qquad (d) \quad P[X \text{ is even}] = \frac{1 - e^{-2}}{2}$$

(a) is still the sum of a geometric progression

(b) follows from the previous result

(c) follows from the previous result taking out a factor of  $1/4$

(d) uses the sum of the two power series of  $e^t$  and  $e^{-t}$ , to knock out the odd terms, evaluated at  $t = 2$ .

2. Let  $Z$  and  $X$  be the numbers of Heads obtained on the first and second tosses respectively. Then the ranges of  $Z$  and  $X$  are both  $\{0, 1, 2, \dots, n\}$ . Now

$$f_X(x) = P[X = x] = \sum_{z=1}^n P[X = x | Z = z] P[Z = z] = \sum_{z=x}^n \binom{z}{x} \left(\frac{1}{2}\right)^z \binom{n}{z} \left(\frac{1}{2}\right)^n$$

using the Theorem of Total probability. Hence

$$f_X(x) = \left(\frac{1}{2}\right)^n \sum_{z=x}^n \frac{z!}{x!(z-x)!} \frac{n!}{z!(n-z)!} \left(\frac{1}{2}\right)^z = \left(\frac{1}{2}\right)^n \binom{n}{x} \sum_{z=x}^n \binom{n-x}{n-z} \left(\frac{1}{2}\right)^z$$

But

$$\sum_{z=x}^n \binom{n-x}{n-z} \left(\frac{1}{2}\right)^z = \sum_{t=0}^m \binom{m}{m-t} \left(\frac{1}{2}\right)^{t+x} = \left(\frac{1}{2}\right)^x \left(1 + \frac{1}{2}\right)^m$$

where  $t = z - x$ , and  $m = n - x$ , using the Binomial Expansion. Hence

$$f_X(x) = \binom{n}{x} \left(\frac{1}{2}\right)^n \left(\frac{1}{2}\right)^x \left(1 + \frac{1}{2}\right)^{n-x} = \binom{n}{x} \frac{3^{n-x}}{2^{2n}} \quad x = 0, 1, 2, \dots, n.$$

Alternately, as all tosses are independent, consider tossing all  $n$  coins twice, and counting the number that show heads twice; this is identical to evaluating  $X$ . Then as each coin shows heads twice with probability  $\left(\frac{1}{2}\right)^2$ ,

$$f_X(x) = \binom{n}{x} \left\{ \left(\frac{1}{2}\right)^2 \right\}^x \left\{ 1 - \left(\frac{1}{2}\right)^2 \right\}^{n-x} = \binom{n}{x} \frac{3^{n-x}}{2^{2n}}$$

as before.

3. Each of the  $n(n+1)/2$  points has equal probability  $p = 2/(n(n+1))$  of being selected. In column  $x$  of the triangular array of points, there are  $x$  points in total; in row  $y$ , there are  $(n+1-y)$  points (for  $x, y = 1, 2, \dots, n$ ) and therefore

$$f_X(x) = P[X = x] = xp = \frac{2x}{n(n+1)} \quad x = 1, 2, \dots, n$$

$$f_Y(y) = P[Y = y] = (n+1-y)p = \frac{2(n+1-y)}{n(n+1)} \quad y = 1, 2, \dots, n$$

4. Can calculate  $F_X$  by integration

$$F_X(x) = \int_{-\infty}^x f_X(t) dt = \int_0^x ct^2(1-t) dt = c \left[ \frac{x^3}{3} - \frac{x^4}{4} \right] \quad 0 < x < 1$$

and  $F_X(1) = 1$  gives  $c = 12$ . Finally,


$$P[X > 1/2] = 1 - P[X \leq 1/2] = 1 - F_X(1/2) = 1 - 12[1/24 - 1/64] = 11/16.$$

5. Valid pdf if (i) it is a non-negative function (that is, if  $k > 0$ ), and (ii) integrates to 1 over the range  $x > 1$ , that is

$$\int_1^{\infty} f_X(x) dx = \int_1^{\infty} \frac{k}{x^{k+1}} dx = \left[ -\frac{1}{x^k} \right]_1^{\infty} = 1 \quad \text{if } k > 0$$

so  $f_X$  is a pdf if  $k > 0$ , and  $F_X(x) = 1 - \frac{1}{x^k}$  for  $x > 1$ .

6. Sketch of  $f_X$ ;



$$F_X(x) = \int_{-\infty}^x f_X(t) dt = \begin{cases} \int_0^x t dt & = \frac{x^2}{2} & 0 < x < 1 \\ \int_0^1 t dt + \int_1^x (2-t) dt & = 2x - \frac{x^2}{2} - 1 & 1 \leq x < 2 \end{cases}$$

Note that  $F_X$  is continuous, and  $F_X(0) = 0$ ,  $F_X(2) = 1$ .

7.  $F_X(1) = 1 \implies \frac{1}{\alpha - \beta}$ , and

$$f_X(x) = \frac{d}{dt} \{F_X(t)\}_{t=x} = \frac{\alpha\beta}{\alpha - \beta} (x^{\beta-1} - x^{\alpha-1}) \quad 0 \leq x \leq 1$$

and zero otherwise, and hence

$$\begin{aligned} E_{f_X} [ X^r ] &= \int_{-\infty}^{\infty} x^r f_X(x) dx = \int_0^1 \frac{\alpha\beta}{\alpha - \beta} x^r (x^{\beta-1} - x^{\alpha-1}) dx \\ &= \frac{\alpha\beta}{\alpha - \beta} \left[ \frac{x^{\beta+r}}{\beta+r} - \frac{x^{\alpha+r}}{\alpha+r} \right]_0^1 \\ &= \frac{\alpha\beta}{(\alpha+r)(\beta+r)} \end{aligned}$$

8. By differentiation,

$$f_X(x) = \frac{d}{dt} \{F_X(t)\}_{t=x} = \frac{2\beta(\beta^2 - x^2)}{(\beta^2 + x^2)^2} \quad 0 \leq x \leq \beta$$

and zero otherwise, and hence

$$\begin{aligned} E_{f_X}[X] &= \int_{-\infty}^{\infty} x f_X(x) dx = \int_0^{\beta} x \frac{2\beta(\beta^2 - x^2)}{(\beta^2 + x^2)^2} dx \\ &= \int_0^{\pi/4} 2\beta^2 \tan \theta \frac{\beta^2(1 - \tan^2 \theta)}{\beta^4(1 + \tan^2 \theta)^2} \beta \sec^2 \theta d\theta \quad (x = \beta \tan \theta) \\ &= 2\beta \int_0^{\pi/4} \tan \theta \frac{(1 - \tan^2 \theta)}{(1 + \tan^2 \theta)} d\theta \\ &= 2\beta \int_0^{\pi/4} \tan \theta \cos 2\theta d\theta \\ &= 2\beta \left[ \frac{1}{2} \tan \theta \sin 2\theta \right]_0^{\pi/4} - \beta \int_0^{\pi/4} \sec^2 \theta \sin 2\theta d\theta \quad (\text{by parts}) \\ &= 2\beta \left[ \frac{1}{2} - \int_0^{\pi/4} \tan \theta d\theta \right] \\ &= 2\beta \left[ \frac{1}{2} - [-\log(\cos \theta)]_0^{\pi/4} \right] \\ &= 2\beta \left[ \frac{1}{2} + \log(\cos \pi/4) \right] = \beta(1 - \log 2) \end{aligned}$$

as  $\cos \pi/4 = 1/\sqrt{2}$ .