

# DYNAMICS NEAR THE PRODUCT OF PLANAR HETEROCLINIC ATTRACTORS

NIKITA AGARWAL, ALEXANDRE RODRIGUES, MICHAEL FIELD

ABSTRACT. Motivated by problems in equivariant dynamics and connection selection in heteroclinic networks, Ashwin and Field investigated the product of planar dynamics where one at least of the factors was a planar homoclinic attractor. However, they were only able to obtain partial results in the case of a product of two planar homoclinic attractors. We give general results for the product of planar homoclinic and heteroclinic attractors. We show that the likely limit set of the basin of attraction of the product of two planar heteroclinic attractors is always the unique one-dimensional heteroclinic network which covers the heteroclinic attractors in the factors. The method we use is general and likely to apply to products of higher dimensional heteroclinic attractors as well as to situations where the product structure is broken but the cycles are preserved.

## 1. INTRODUCTION

One way of analyzing the dynamics of coupled dynamical systems is to first understand the dynamics of product systems (that is, uncoupled systems) and then perturb by adding coupling. In the case of two uncoupled dynamical systems, the perturbation could be to a skew product system where the base system weakly forces the second system. Motivated by problems in equivariant dynamics and coupled systems, Ashwin and Field [4] made a preliminary study of product dynamics when one of the factors was a (planar) homoclinic attractor. Strong results were obtained when the other factor was an attracting limit cycle — perhaps not so surprisingly, the main result (that the product was a minimal Milnor attractor [4, Theorem 1.2]) depended on non-trivial results from metric number theory. Strong results were also

---

*Date:* July 9, 2011.

Research of NA, MF and AR supported in part by NSF Grant DMS-0806321, research of AR supported in part by Centro de Matemática da Universidade do Porto (CMUP) and Fundação para a Ciência e Tecnologia (FCT), Portugal, through the programs POCTI and POSI with European Union and national fundings and also by the grant SFRH/BD/28936/2006 of FCT.

found when the second factor was a basic hyperbolic set. However, if both factors were planar homoclinic attractors, results were only obtained for a very restricted model [4, §6, Theorem 6.13].

In this paper we give quite general results about the *likely limit sets* of products of two-dimensional homoclinic and heteroclinic attractors. Although we restrict to products of two-dimensional systems, we believe that the methods we use have much wider applicability both to higher dimensional systems and to systems which are perturbations of a product.

Before stating our main result and giving examples, we need to recall the definition of the ‘likely limit set’ (details and references are given in the next section). Let  $M$  be a compact manifold, possibly with boundary, with measure  $\ell$  which we assume is locally equivalent to Lebesgue measure. Suppose that  $X$  is a compact indecomposable attractor for the flow  $\Phi_t : M \rightarrow M$  and that  $X$  has basin of attraction  $\mathcal{B}(X)$  which we assume has strictly positive measure. It may be the case that  $\ell$ -almost all points in  $\mathcal{B}(X)$  are forward asymptotic to a proper subset of  $X$ . We capture this idea by defining the likely limit set of  $\mathcal{B}(X)$  to be the smallest compact flow invariant subset  $Z$  of  $X$  with the property that for  $\ell$ -almost all points  $x \in \mathcal{B}(X)$ , the omega limit set  $\omega(x) \subset Z$ .

Suppose that  $\Phi_t$  is a  $C^1$  flow on the compact surface  $M$ , possibly with boundary. A *heteroclinic network* for the flow  $\phi_t$  consists of a closed connected 1-dimensional  $\Phi_t$ -invariant subset  $\Sigma$  of  $M$  which is the union of a finite set  $\mathcal{E}(\Sigma) = \mathcal{E}$  of hyperbolic saddle points and a finite set of  $\Phi_t$ -trajectories connecting equilibria in  $\mathcal{E}$  such that the graph defined by equilibria (vertices) and trajectories (edges) is strongly connected (given two equilibria  $\mathbf{p}, \mathbf{q} \in \mathcal{E}$ , there exists a finite chain of trajectories in  $\Sigma$  joining  $\mathbf{p}$  to  $\mathbf{q}$ ). The simplest example of a heteroclinic network is a homoclinic loop consisting of one equilibrium and one trajectory (see below). We say that the heteroclinic network  $\Sigma$  is a *heteroclinic attractor* if the basin of attraction of  $\Sigma$  is a neighbourhood of  $\Sigma$  in  $M$  (below we allow for one-sided attractors, such as a homoclinic loop).

**Theorem 1.1.** *Let  $M_1, M_2$  be compact surfaces and  $\Sigma_1 \subset M_1, \Sigma_2 \subset M_2$  be heteroclinic attractors for  $C^2$  flows  $\phi_t^i : M_i \rightarrow M_i$ ,  $i = 1, 2$ . The likely limit set of  $\mathcal{B}(\Sigma_1 \times \Sigma_2)$  for the product flow  $\phi_t^1 \times \phi_t^2$  is the heteroclinic network*

$$(\Sigma_1 \times \mathcal{E}(\Sigma_2)) \cup (\mathcal{E}(\Sigma_1) \times \Sigma_2) \subset M_1 \times M_2.$$

In figure 1 we show a number of examples of homoclinic and heteroclinic attractors.

The attracting planar homoclinic loop  $\Sigma \subset D$  shown in figure 1(a) is the simplest example of a planar homoclinic attractor. In this case the




FIGURE 1. Homoclinic attractors on surfaces. (a) One sided homoclinic attractor, (b) figure-of-eight homoclinic attractor, (c) heteroclinic attractor on the Klein bottle, (d) heteroclinic attractor on projective space

single equilibrium point  $\mathbf{p}$  on the loop is a saddle point with contraction dominating the expansion. This loop is a one-sided attractor — nothing is said about the dynamics on the complement of the region  $D$  enclosed by the loop. If we take two such attracting planar homoclinic loops  $\Sigma_1 \subset D_1$ ,  $\Sigma_2 \subset D_2$ , with corresponding equilibria  $\mathbf{p}_1$ ,  $\mathbf{p}_2$ , then it is a consequence of results in [4] that the only possibilities for the likely limit set of  $\mathcal{B}(\Sigma_1 \times \Sigma_2) \subset D_1 \times D_2$  are either  $\Sigma_1 \times \Sigma_2$  or the homoclinic network  $(\Sigma_1 \times \{\mathbf{p}_2\}) \cup (\{\mathbf{p}_1\} \times \Sigma_2)$ . It follows from theorem 1.1 that the second case is the only possibility.

The figure of eight homoclinic attractor  $\Sigma$  shown in figure 1(b) is attracting on both sides provided that the single equilibrium point  $\mathbf{p}$  on the loop is a saddle point with contraction dominating the expansion. We consider the product of two loops of this type or the product with a one-sided homoclinic attractor of the type shown in figure 1(a). For example, if we take the product of figure of eight loops  $\Sigma_1$ ,  $\Sigma_2$ , then theorem 1.1 implies that the likely limit set of  $\mathcal{B}(\Sigma_1 \times \Sigma_2)$  is always the homoclinic network  $(\Sigma_1 \times \{\mathbf{p}_2\}) \cup (\{\mathbf{p}_1\} \times \Sigma_2)$ . In this case, there

is a richer structure present as if we look at the product of  $\mathcal{B}(\Sigma_1^u)$  (see figure 1) with that part of the basin of attraction of  $\Sigma_2$  exterior to the loop, then the likely limit set is  $(\Sigma_1^u \times \{\mathbf{p}_2\}) \cup (\{\mathbf{p}_1\} \times \Sigma_2)$ .

In figure 1(c,d), we show two simple examples of heteroclinic attractors defined on the Klein bottle (c) and 2-dimensional real projective space (d). In both figures we identify opposite edges of the square so that  $\alpha$  is identified with  $\bar{\alpha}$ ,  $\beta$  with  $\bar{\beta}$ . The vector field on the Klein bottle has two saddle point equilibria  $\mathbf{p}, \mathbf{q}$  on the heteroclinic network  $\Sigma = \Sigma^a \cup \Sigma^b$ , where in figure 1(c) we have indicated the unique edge of the heteroclinic cycle  $\Sigma^b$  (respectively,  $\Sigma^a$ ), which is not common to  $\Sigma^a$  (respectively,  $\Sigma^b$ ).

The vector field on projective space also has two saddle point equilibria  $\mathbf{p}, \mathbf{q}$  on the heteroclinic network  $\Sigma = \Sigma^a \cup \Sigma^b$ . In this case the complement of the network has *three* connected regions.

If we take the product of the heteroclinic attractor  $\Sigma_1$  on the Klein bottle with that the heteroclinic attractor  $\Sigma_2$  on projective space, then it follows from theorem 1.1 that the likely limit set of  $\mathcal{B}(\Sigma_1 \times \Sigma_2)$  is the heteroclinic network  $\Sigma = (\Sigma_1 \times \{\mathbf{p}_2, \mathbf{q}_2\}) \cup (\{\mathbf{p}_1, \mathbf{q}_1\} \times \Sigma_2)$ .

*Remarks 1.2.* (1) For general attracting heteroclinic networks, we expect the presence of *essentially asymptotically stable* subnetworks (see Brannath [6] and Melbourne [19]). More precisely, if  $\Sigma$  is an attracting heteroclinic network associated to a finite number of equilibria with only real eigenvalues, then the connections associated to the strongest expanding eigenvalues determine a possibly smaller attractor. If the attractor is a heteroclinic cycle then it is essentially asymptotically stable (in a neighbourhood of the network, almost all orbits converge to the cycle). An explicit example of this phenomenon was given by Kirk and Silber [16]. Given an asymptotically stable heteroclinic network  $\Sigma$ , define the principal out-connection of a saddle to be the heteroclinic connection corresponding to the most positive expanding eigenvalue of the linearization at the node. Ashwin and Chossat [5] conjectured the existence of an essentially asymptotically stable subnetwork  $\Sigma^* \subset \Sigma$  containing the principal connections that forms part of the attractor  $\Sigma$ . A general proof of essential asymptotic stability can be based on the Strong Lambda Lemma of Deng [8] but this result lies beyond the scope of the present paper. By contrast, products of heteroclinic attractors generally lead to heteroclinic networks without proper essentially asymptotically stable subnetworks. This is irrespective of the relative strength of eigenvalues at the equilibria. In all of the examples described above, the likely limit sets contain no proper essentially asymptotically stable heteroclinic cycles.

(2) For a product of planar heteroclinic cycles, the invariant subspaces forced by the product structure are least codimension two. In the absence of proper essentially asymptotically stable cycles, it is natural to ask whether switching (in the sense described by Aguiar *et al.* [2] and Homburg *et al.* [13]) can occur. We briefly address the study of this phenomenon at the end of section 3.

At the suggestion of a referee, we include some heuristic comments about the main result.




FIGURE 2. A product of homoclinic attractors

Suppose that the planar flows  $\phi_t^1, \phi_t^2$  have homoclinic attractors  $\Sigma_1, \Sigma_2$  respectively. Consider a segment  $\gamma_1$  of a trajectory of  $\phi_t^1$ , with initial condition  $x \notin \Sigma_1$ , that starts at  $a \in A$  and makes one circuit of the loop, arriving back at  $c$  (see figure 2). Choose times  $t_a, t_b, t_c$  such that  $a = \phi_{t_a}^1(x), b = \phi_{t_b}^1(x), c = \phi_{t_c}^1(x)$ . If we assume the flow is linear on  $A$ , then it is easy to show that  $t_c - t_b = O(|\log a|)$  and so  $t_b - t_a \rightarrow \infty$  as  $a \rightarrow 0$ . On the other hand  $t_c - t_b$  will be uniformly bounded above. If we pick a random time  $t \in [t_a, t_c]$ , then the probability  $p$  that  $\phi_t^1(x) \notin A$  is  $(t_c - t_b)/(t_c - t_a)$  and so  $p \rightarrow 0$  as  $a \rightarrow 0$ . Similar comments hold for circuits  $\gamma_2$  of the second system. If we assume the events of lying outside  $A$  and  $B$  are independent, then it follows that for almost all initial conditions  $(x, y)$ ,  $\omega(x, y)$  contains no points of  $(\Sigma_1 \setminus A) \times (\Sigma_2 \setminus B)$ . The required result on the likely limit set would then follow by letting the diameters of  $A$  and  $B$  decrease to zero. However, since we are dealing with a deterministic system, we cannot assume independence of events (though we can expect a ‘decay of correlations’). In practice, what we have to do is show that for almost all initial conditions  $(x, y)$ , we can find a future time  $T = T(x, y)$  such that for  $t \geq T$ , either  $\phi_t^1(x) \in A$  or  $\phi_t^2(y) \in B$ . This in turn implies that for almost all initial conditions

$(x, y)$ ,  $\omega(x, y)$  contains no points of  $(\Sigma_1 \setminus A) \times (\Sigma_2 \setminus B)$ . Our basic argument depends on obtaining precise estimates on the times  $t$  for which we can have  $\phi_t^1(x) \notin A$  and  $\phi_t^2(y) \notin B$  and then applying the Borel-Cantelli lemma (the part that does not require independence of events!) The analysis is fairly straightforward if we assume linear connection maps. However, the general case is tricky and does *not* follow directly from the special case. This, in itself, is a caution about making linearity assumptions on connection maps in heteroclinic networks: the nonlinearities are highly significant. To obtain our general results, we require flows to be at least  $C^2$ . *Our methods fail completely if the flows are only  $C^1$  (or  $C^{1+\varepsilon}$ ) even if we assume  $C^1$  linearizability at the equilibria.* Indeed, it is quite possible that our results do not hold if the flows are only  $C^1$ .

We describe the contents of the paper by section. In section 2, we review basic definitions and results on Milnor attractors and the likely limit set. We also establish some conventions for our subsequent analysis of planar and surface attracting homoclinic and heteroclinic cycles. In section 3, we establish notational conventions and outline the strategy of the proof of the main result for the case of the product of two planar attracting homoclinic loops. The key and new ingredient is the use of the Borel-Cantelli lemma. In section 4 we give the proof for the product of two planar attracting homoclinic loops subject only to a restriction on the connection maps — we assume they are linear. In section 5, we show how our methods easily extend to general products of heteroclinic attractors, including figure eight homoclinic cycles and attracting heteroclinic cycles. We continue to assume the linearity restriction on connection maps. In section 6, we remove the linearity assumption on connection maps. The resulting analysis is delicate, especially in the resonant case where we assume that both heteroclinic cycles have the same asymptotic attractivity (for a product of homoclinic loops, this amounts to the ratio of the eigenvalues at the equilibria being equal). We present the details only in the case of a product of attracting homoclinic loops but the extension of our methods and results to the general case is clear. At first reading, we would advise (and expect) most readers to omit the details in section 6. Overall these three sections illustrate how we use our argument based on the Borel-Cantelli lemma in a technically simple situation (section 4); how we extend to more general networks (section 5) and finally how we handle the arguments in the technically more demanding case when we make no simplifying assumptions on connection maps (section 6).

In sections 7 and 8, we show the results of numerical simulations of product systems as well as consider cases where we break homoclinic

connections but preserve the product structure. We conclude with a brief discussion of possible generalizations and extensions of our results.

## 2. PRELIMINARIES

**2.1. Milnor attractors and the likely limit set.** Let  $M$  be a differential manifold, possibly with boundary, and let  $\ell$  denote a measure on  $M$  locally equivalent to the Lebesgue measure on charts (for example, if  $M$  is an orientable Riemannian manifold, then  $\ell$  can be the measure defined by the Riemannian volume form). If  $Z$  is a measurable subset of  $M$  with  $\ell(Z) \neq 0$ , we let  $\mathcal{F}(Z)$  denote the set of measurable subsets  $Z'$  of  $Z$  such that  $\ell(Z \setminus Z') = 0$ .

Suppose that  $\Phi_t : M \rightarrow M$  is a  $C^1$  flow (or semi-flow) on  $M$ . Given  $x \in M$ , let

$$\omega(x) = \bigcap_{T>0} \overline{\{\Phi_t(x) \mid t \geq T\}}$$

denote the  $\omega$ -limit set of the trajectory through  $x$ .

If  $X$  is a compact invariant subset of  $M$ , we let  $\mathcal{B}(X) = \{x \in M \mid \omega(x) \subset X\}$  denote the *basin of attraction* of  $X$ . We recall the definitions of a Milnor attractor and minimal Milnor attractor (for more details we refer to Milnor [20]).

**Definition 2.1** (Milnor [20]). A compact invariant subset  $X$  of  $M$  is a *Milnor attractor* if

- (1)  $\ell(\mathcal{B}(X)) > 0$ ;
- (2) for any proper compact invariant subset  $Y$  of  $X$ ,  $\ell(\mathcal{B}(X) \setminus \mathcal{B}(Y)) > 0$ .

We say  $X$  is a *minimal (Milnor) attractor* if for all proper compact invariant subsets  $Y$  of  $X$ ,  $\ell(\mathcal{B}(Y)) = 0$ .

*Remark 2.2.* A Milnor attractor  $X$  is minimal iff there is a full measure subset  $B$  of  $\mathcal{B}(X)$  such that  $\omega(x) = X$  for all  $x \in B$ .

**Definition 2.3** (cf. Milnor [20]). Let  $Z \subset M$  be measurable with  $\ell(Z) > 0$ , and  $Z$  forward  $\Phi_t$ -invariant. The *likely limit set*  $\mathcal{L}(Z)$  of  $Z$  is the smallest closed  $\Phi_t$ -invariant subset of  $Z$  that contains all  $\omega$ -limit sets except for a subset of  $Z$  of zero measure. That is,

$$\mathcal{L}(Z) = \bigcap_{Z' \in \mathcal{F}(Z)} \overline{\{\omega(x) \mid x \in Z'\}}.$$

*Remarks 2.4.* (1) If  $Z$  is relatively compact, then  $\mathcal{L}(Z)$  is a non-empty, compact  $\Phi_t$ -invariant subset of  $M$ .

(2) The definition of the likely limit set applies to measurable subsets  $Z \subset M$  with  $\ell(Z) > 0$  that are not necessarily forward  $\Phi_t$ -invariant: since the flow is assumed  $C^1$ , it preserves measure zero sets and from

this it follows straightforwardly that  $\mathcal{L}(Z) = \mathcal{L}(\cup_{t \geq 0} \Phi_t(Z))$ . (Of course,  $\mathcal{L}(Z)$  may be empty if  $\cup_{t \geq 0} \Phi_t(Z)$  is not relatively compact.)

We recall two results from [4] about likely limit sets.

**Lemma 2.5** (Ashwin & Field [4, Lemma 2.3]). *Let  $Z \subset M$  be measurable with  $\ell(Z) > 0$ , and  $Z$  forward  $\Phi_t$ -invariant.*

- (1)  $x \in \mathcal{L}(Z)$  iff for all  $\varepsilon > 0$  and all  $Z' \in \mathcal{F}(Z)$  there exists  $a \in Z'$  such that  $B_\varepsilon(x) \cap \omega(a) \neq \emptyset$  ( $B_\varepsilon(x)$  denotes the  $\varepsilon$  ball about  $x$ );
- (2)  $\mathcal{L}(Z)$  is a minimal Milnor attractor iff for all  $x \in \mathcal{L}(Z)$  and all  $\varepsilon > 0$  and all  $H \subset Z$  with  $\ell(H) > 0$ , we have

$$\ell(\{a \in H \mid B_\varepsilon(x) \cap \omega(a) \neq \emptyset\}) > 0.$$

**Theorem 2.6** (cf. Ashwin & Field [4, Theorem 1.1]). *Let  $M_1, M_2$  be compact manifolds, possibly with boundary and  $\Phi_t = (\phi_t^1, \phi_t^2)$  be a product of  $C^1$  flows on  $M_1 \times M_2$ . Suppose that  $X_i \subset M_i$  are forward  $\phi_t^i$ -invariant measurable subsets of  $M_i$  of strictly positive measure,  $i = 1, 2$ . Then  $\mathcal{L}(X_1 \times X_2)$  is invariant under the  $\mathbb{R}^2$  action defined by  $(\phi_t^1, \phi_s^2)$ ,  $t, s \in \mathbb{R}^2$ .*

*Proof.* It is no loss of generality to assume that the  $X_i$  are  $\phi_t^i$ -invariant subsets of  $M_i$  (replace  $X_i$  by  $\cup_{t \in \mathbb{R}} \phi_t^i(X_i)$ ). The proof given in [4] for the case  $X_i = M_i$ ,  $i = 1, 2$ , extends trivially to the case where the  $X_i \subset M_i$  are  $\phi_t^i$ -invariant measurable subsets of  $M_i$  of strictly positive measure – the crucial point is that the flows  $\phi_t^i$  preserve sets of measure zero.  $\square$

**2.2. Planar homoclinic attractors.** Suppose that we are given a  $C^2$  semi-flow  $\Phi_t$  defined on some region  $D^*$  of  $\mathbb{R}^2$  containing the origin. We assume that the origin is a hyperbolic saddle with associated eigenvalues  $-\mu < 0 < \lambda$  and eigen-directions the  $x$ - and  $y$ -axes of  $\mathbb{R}^2$  respectively. We also assume that there is a homoclinic cycle  $\Sigma \subset D^*$  connecting the origin (see figure 3).

If  $\mu > \lambda$ , then it is well known that  $\Sigma$  is a (one-sided) attracting homoclinic cycle. More precisely, if we let  $D \subset D^*$  be the compact region in  $\mathbb{R}^2$  with boundary  $\Sigma$ , then there exists an open neighbourhood  $N$  of  $\Sigma$  in  $D$  such that for all  $x \in N \setminus \Sigma$ ,  $\omega(x) = \Sigma$ . We may choose  $N$  so that  $N$  is forward  $\Phi_t$ -invariant with smooth interior boundary  $\partial N$  and such that the trajectories of  $\Phi_t$  intersect  $\partial N$  transversally (see figure 4 and note that if  $x \notin D$ , then  $\omega(x) \not\subset \Sigma$ ).

Given the setup shown in figure 4,  $N$  is a subset of the basin of attraction  $\mathcal{B}(\Sigma)$  of  $\Sigma$  and, of course,  $\mathcal{L}(N) = \mathcal{L}(\mathcal{B}(\Sigma)) = \Sigma$ . In future, we always assume that a homoclinic cycle  $\Sigma$  of this type is the boundary of a compact region  $D \subset \mathbb{R}^2$  and comes with a choice of neighbourhood  $N$




FIGURE 3. Attracting homoclinic loop in region  $D^* \subset \mathbb{R}^2$ . Inside the region bounded by the homoclinic cycle  $\Sigma$ , trajectories approach the cycle. Outside this region, trajectories eventually go away from the cycle.




FIGURE 4. Interior neighbourhood  $N$  of  $\Sigma$ . Note that  $N$  is constructed in such a way that it is forward flow invariant and trajectories of  $\Phi_t$  intersect  $\partial N$  transversely.

satisfying the properties described above. We denote by the quadruple  $(\Sigma, \Phi_t, D, N)$ .

*Remark 2.7.* Noting remarks 2.4(2), if  $A$  is an open interior neighbourhood of  $(0, 0)$  (see figure 4), then  $\mathcal{L}(A) = \Sigma$ .

We may carry out a similar process for an attracting ‘figure of eight’ homoclinic cycle in the plane or, more generally an attracting planar heteroclinic cycle such as the Guckenheimer-Holmes cycle [11], [10, §5.2] or the heteroclinic cycles on the Klein bottle and projective space shown in figure 1.

### 3. PRODUCTS OF PLANAR ATTRACTING HOMOCLINIC LOOPS

For the next two sections we assume that  $(\Sigma_i, \phi_t^i, D_i, N_i)$ ,  $i = 1, 2$ , are planar attracting homoclinic loops (‘homoclinic attractors’). Recall that we assume  $\partial D_i = \Sigma_i$ ,  $\phi_t^i$  is a  $C^2$  flow on  $D_i$ , and  $N_i \subset \mathcal{B}(\Sigma_i)$  is a forward  $\phi_t^i$ -invariant open interior neighbourhood of  $\Sigma_i$ . We assume that both homoclinic attractors have a saddle point at  $(0, 0)$ , and corresponding eigenvalues  $-\mu_i < 0 < \lambda_i$ , where  $\rho_i = \mu_i/\lambda_i > 1$ ,  $i = 1, 2$  (in the terminology of L Shilnikov, the  $\rho_i$  are the *saddle indices* of the homoclinic loops [22, Chapter 13]). Set  $\Sigma = (\Sigma_1 \times \{(0, 0)\}) \cup (\{0, 0\} \times \Sigma_2) \subset D_1 \times D_2 \subset \mathbb{R}^4$  and let  $\mathbf{0} = (0, 0, 0, 0) \in \Sigma$  denote the unique equilibrium for the product flow in  $N_1 \times N_2$ . Let  $\Phi_t = (\phi_t^1, \phi_t^2)$  denote the product flow on  $D_1 \times D_2$ .

Using theorem 2.6, it was shown in Ashwin & Field [4, Theorem 6.1] that either  $\mathcal{L}(N_1 \times N_2) = \Sigma$  or  $\mathcal{L}(N_1 \times N_2) = \Sigma_1 \times \Sigma_2$ . Our aim is to prove that we always have  $\mathcal{L}(N_1 \times N_2) = \Sigma$ . For the remainder of this section, we outline the strategy that we use for the proof as well as establish notational conventions.




FIGURE 5. Linearizing coordinates at the origin.

Since  $(0, 0)$  is a hyperbolic saddle point of a  $C^2$  planar flow, it follows by a result of Hartman [12] that we can always choose a  $C^1$ -linearization of  $\phi_t^i$  on some closed neighbourhood  $A_i$  of  $(0, 0)$ ,  $i = 1, 2$  (the  $C^2$  regularity of the flow will play a key role in section 6). Linearly rescaling coordinates, we may assume that  $A_i = [0, 1] \times [0, 1] \subset N$ . Let  $A_i^\circ = [0, 1) \times [0, 1)$  denote the interior of  $A_i$  in  $D_i$ . We may choose the  $A_i$  so that the forward  $\phi_t^i$ -trajectory through  $(1, 1)$  meets  $\{1\} \times [0, 1] \subset \partial A_i$  after one circuit of the loop (see figure 5).

Given  $i \in \{1, 2\}$ , let  $U_i = \{1\} \times [0, 1]$ ,  $V_i = [0, 1] \times \{1\} \subset \partial A_i$  denote the vertical and horizontal interior boundaries of  $A_i$  (see figure 6).

We adopt the convention that if  $y \in [0, 1] \approx U_i$ , then  $\phi_t^i(y)$  is defined to be  $\phi_t^i(1, y)$ . We similarly identify a point  $(x, 1) \in V_i$  with  $x \in [0, 1]$ .

We have a  $C^1$  time of first return map  $T_i : V_i \rightarrow \mathbb{R}$  and associated  $C^1$  connection map  $C_i : V_i \rightarrow U_i$  defined by  $C_i(x) = \phi_{T_i(x)}^i(x)$ ,  $x \in V_i$ .

Let  $\tau_i^+ = \sup_{x \in V_i} T_i(x)$ ,  $\tau_i^- = \inf_{x \in V_i} T_i(x)$ . Obviously,  $\tau_i^+ \geq \tau_i^- > 0$ . Since the time it takes for a trajectory starting at  $y \in U_i$  to exit  $A_i$  through  $V_i$  grows without bound as  $y \rightarrow 0^+$ , it is easy to verify that every trajectory in  $N_1 \times N_2$  passes through  $A_1 \times A_2$ . From this it follows that  $\mathcal{L}(A_1 \times A_2) = \mathcal{L}(N_1 \times N_2)$ .




FIGURE 6. Notation and setup for the sets  $A_1, A_2$ : all trajectories with initial conditions on  $U_i \setminus \{(1, 1)\}$  (respectively  $V_i \setminus \{(1, 1)\}$ ), are inside  $A_i$  for small strictly positive (respectively, negative) time.

Given  $b \in U_1$ ,  $b > 0$ , let  $a < b$  denote the first point of intersection of the forward  $\phi_t^1$ -trajectory through  $b$  with  $U_1$ . Similarly, given  $d \in U_2$ ,  $d > 0$ , define  $c < d$  to be the first point of intersection of the forward  $\phi_t^2$ -trajectory through  $d$  with  $U_2$ . (See figure 6.)

Given  $y \in [a, b]$ , let  $0 < s_1^y(1) < s_1^y(2) < \dots$  denote the sequence of positive times  $t$  at which  $\phi_t^1(y) \in V_1$  and  $0 = t_1^y(0) < t_1^y(1) < t_1^y(2) < \dots$  denote the sequence of positive times  $t$  at which  $\phi_t^1(y) \in U_1$ . Observe that  $t_1^y(n) - s_1^y(n) \in [\tau_1^-, \tau_1^+]$ .

Let  $(x_n^1) \subset V_1$  be the decreasing sequence of points of intersection of the forward trajectory through  $y \in [a, b]$  with  $V_1$  and  $(y_n^1)$  be the corresponding sequence of points of intersection with  $U_1$ . We have

$$\begin{aligned} y_n^1 &= \phi_{T_1(x_n^1)}^1(x_n^1), \\ x_n^1 &= \phi_{s_1^y(n)}^1(y) = \phi_{s_1^y(n) - s_1^y(1)}^1(x_1^1), \\ y_n^1 &= \phi_{t_1^y(n)}^1(y) = \phi_{s_1^y(n) + T_1(x_n^1)}^1(y), \\ s_1^y(n+1) - t_1^y(n) &= \lambda_1^{-1} \log(y_n^1)^{-1}, \\ x_{n+1}^1 &= (y_n^1)^{\mu_1/\lambda_1} = (y_n^1)^{\rho_1}, \end{aligned}$$

where the last two statements use the linearity of the flow  $\phi_t^1$  on  $A_1$  and hold for  $n \geq 0$ . Similar statements hold for  $\phi_t^2$ .

**Definition 3.1.** Given  $n \in \mathbb{N}$ ,  $y \in [a, b]$ , we say  $t > 0$  is  $n$ -singular for  $A_1$  if  $t \in [s_1^y(n), t_1^y(n)]$ . If  $t \geq s_1^y(n)$ , we say the trajectory  $\phi_t^1(y)$  has made at least  $n$ -turns about the homoclinic cycle  $\Sigma_1$ .

*Remarks 3.2.* (1) If  $t$  is  $n$ -singular for  $A_1$ , then  $\phi_t^1(y) \notin A_1^\circ$ .  
(2) The definition of making at least  $n$ -terms about  $\Sigma_1$  implicitly depends on  $A_1$ .

Let  $\mathcal{C} \subset \Sigma_1 \times \Sigma_2$  consist of all pairs  $(p, q) \in \Sigma_1 \times \Sigma_2$  such that  $p, q \notin [0, 1) \times \{0\} \cup \{0\} \times [0, 1)$ . If we let  $\kappa_i \subset \Sigma_i$  be the closed arcs defined by  $\kappa_i = \Sigma_i \setminus A_i^\circ$ , then  $\mathcal{C} = \kappa_1 \times \kappa_2$ .

**Definition 3.3.** Given  $A_1, A_2$  as defined above, a point  $(X, Y) \in A_1 \times A_2$  is a *bad point* if  $\omega(X, Y) \cap \mathcal{C}^\circ \neq \emptyset$  ( $\mathcal{C}^\circ$  denotes the interior of  $\mathcal{C}$  in  $\Sigma_1 \times \Sigma_2$ ).

*Remark 3.4.* We shall show that the set of bad points in  $A_1 \times A_2$  has measure zero. It then follows immediately from theorem 2.6 that  $\mathcal{L}(A_1 \times A_2) = \Sigma$ .

Let  $E = [a, b] \times [c, d] \subset U_1 \times U_2$ . For  $n \in \mathbb{N}$ , define  $E_n \subset E$  by

$$E_n = \{(y, z) \in E \mid \exists t \in [s_1^y(n), t_1^y(n)] \text{ such that } \phi_t^2(z) \notin A_2^\circ\}.$$

We refer to points of  $E_n$  as the  $n$ -bad points of  $E$ . Observe that  $(y, z) \in E_n$  if and only if there exists  $t > 0$ ,  $m \in \mathbb{N}$  such that  $t$  is  $n$ -singular for  $A_1$  and  $m$ -singular for  $A_2$ . In particular,  $\phi_t^2(z) \notin A_2^\circ$ . If, for  $m \in \mathbb{N}$ , we define

$$E_{n,m} = \{(y, z) \in E \mid [s_1^y(n), t_1^y(n)] \cap [s_2^z(m), t_2^z(m)] \neq \emptyset\},$$

then  $E_n = \cup_{m \geq 1} E_{n,m}$ .

**Lemma 3.5.** *If  $(y, z) \in E$ , then a point  $(p, q) \in \mathcal{C}^\circ \cap \omega(y, z)$  only if there exists an infinite increasing sequence  $n_1 < n_2 < \dots$  such that  $(y, z) \in \cap_{j \geq 1} E_{n_j}$ . Conversely, if there exists an infinite increasing sequence  $n_1 < n_2 < \dots$  such that  $(y, z) \in \cap_{j \geq 1} E_{n_j}$ , then  $(p, q) \in \mathcal{C} \cap \omega(y, z)$ .*

*Proof.* If there exists  $N \in \mathbb{N}$  such that  $(y, z) \notin E_n$ ,  $n \geq N$ , then all the limit points of the  $\Phi_t$ -trajectory through  $(y, z)$  must lie in  $(A_1 \times \Sigma_2) \cup (\Sigma_1 \times A_2)$  which is disjoint from  $\mathcal{C}^\circ$ . Conversely, suppose  $(y, z) \in \cap_{j \geq 1} E_{n_j}$ . Using the compactness of  $\Sigma_1 \times \Sigma_2$ , we can pick a subsequence  $(m_j)$  of  $(n_j)$  and sequence  $t_{m_1} < t_{m_2} < \dots$  with  $t_{m_j} \in [s_1^y(m_j), t_1^y(m_j)]$  and  $\phi_{t_{m_j}}^2(z) \notin A_2^\circ$ , such that  $(\Phi_{t_{m_j}}(y, z))$  converges to a point in  $\mathcal{C}$ .  $\square$

*Applying the Borel-Cantelli lemma.* Let  $E_\infty$  denote the subset of  $E$  consisting of points  $(y, z)$  such that there exists an infinite increasing sequence  $(n_j)$  such that  $(y, z) \in \cap_{j \geq 1} E_{n_j}$ . We have  $E_\infty = \cap_{N \geq 1} \cup_{n \geq N} E_n$ . It follows from lemma 3.5 that the bad points of  $E$  are a subset of  $E_\infty$ . Take Lebesgue measure  $\ell_2$  on  $E$ . Our main work will be to show  $\sum_{n=1}^\infty \ell_2(E_n) < \infty$ . It then follows from the Borel-Cantelli lemma that  $\ell_2(E_\infty) = 0$  and so there is a full measure subset  $E'$  of  $E$  such that  $\omega(y, z) \cap \mathcal{C}^\circ = \emptyset$  for all  $(y, z) \in E'$ .

The proof we give for the convergence of  $\sum_{n=1}^\infty \ell_2(E_n)$  applies to all products  $E = [a, b] \times [c, d] \subset (0, 1] \times (0, 1] \subset U_1 \times U_2$ , where  $a, c$  are the points of first return. Since  $(0, 1] \times (0, 1]$  can be written as a countable union of such products, it follows by the  $\sigma$ -additivity of  $\ell_2$  that the set of bad points in  $U_1 \times U_2$  has zero measure.

Furthermore, we are able to show that our arguments do not depend on the particular choice of linearizing neighbourhood  $A_1 \times A_2$ : we can replace  $E = (\{1\} \times [a, b]) \times (\{1\} \times [c, d])$  by any product of vertical intervals from  $A_1 \times A_2$ . That is, suppose  $x, x', b, d \in (0, 1]$ , and define corresponding intervals  $I_x = \{x\} \times [a, b] \subset A_1$ ,  $I_{x'} = \{x'\} \times [c, d] \subset A_2$  such that  $a$  is the first point of intersection of the forward  $\phi_t^1$ -orbit through  $(x, b)$  with  $\{x\} \times [0, 1]$ , and similarly for  $c$ . Set  $E = I_x \times I_{x'}$ . Now linearly rescale the horizontal coordinates to obtain a new box linearizing neighbourhood  $A_1^* \times A_2^* \subset A_1 \times A_2$  such that in the new coordinates  $A_1^*, A_2^* = [0, 1] \times [0, 1]$  and  $I_x = [a, b] \subset \{1\} \times (0, 1]$ ,  $I_{x'} = [c, d] \subset \{1\} \times (0, 1]$  (note that the vertical coordinates are unchanged). By the original argument, it follows there is a full measure subset  $E'$  of  $E$  such that  $\omega(y, z) \cap \mathcal{C}^\circ = \emptyset$  for all  $(y, z) \in E'$  ( $\mathcal{C}$  is expanded when we use the smaller box  $A_1^* \times A_2^*$  so we can safely take the *original*  $\mathcal{C}$  for this argument). Using the argument of the previous paragraph,

we deduce that the bad points are a measure zero subset of  $(\{x\} \times (0, 1]) \times (\{x'\} \times [0, 1]) \subset A_1 \times A_2$  for all  $x, x' \in (0, 1]$ . Since the set of bad points in  $A_1 \times A_2$  is easily seen to be measurable, it follows by Fubini's theorem that the set of bad points form a measure zero subset of  $A_1 \times A_2$ . Therefore there is a full-measure subset  $A'$  of  $A_1 \times A_2$  for which  $\omega(y, z) \cap \mathcal{C}^\circ = \emptyset$  for all  $(y, z) \in A'$  proving that  $\mathcal{L}(A_1 \times A_2) \neq \Sigma_1 \times \Sigma_2$ . Hence,  $\mathcal{L}(N_1 \times N_2) = \Sigma$ .

*Remark 3.6.* The final step uses the  $\mathbb{R}^2$ -invariance of the likely limit set (theorem 2.6). However, we could have avoided this by observing that our arguments show the existence of a full-measure subset  $A'$  of  $A_1 \times A_2$  for which  $\omega(y, z) \cap \mathcal{C}^\circ = \emptyset$  for all  $(y, z) \in A'$ . Since this holds for a base of box neighbourhoods  $A_1 \times A_2$  of  $\mathbf{0}$ , we deduce that  $\mathcal{L}(N_1 \times N_2) \subset \Sigma$ . Of course, for product dynamics we can infer equality. However, in other situations where the product structure is broken but the attracting homoclinic loops persist, the inclusion  $\mathcal{L}(N_1 \times N_2) \subset \Sigma$  may be strict (this is of particular interest for products of planar 'figure of eight' attracting homoclinic loops.)

#### 4. PRODUCTS OF HOMOCLINIC LOOPS: PROOF OF A SPECIAL CASE

In this section, we present the proof for products of planar attracting homoclinic loops under a simplifying assumption on the connection maps  $C_i : V_i \rightarrow U_i$ . We assume that for both flows  $\phi_t^i$ , there exist strictly positive constants  $\tau_1, \tau_2, m_1, m_2$  such that

$$(4.1) \quad T_i(x) = \tau_i, \quad C_i(x) = m_i x, \quad x \in U_i.$$

We give the proof for general connection maps in section 6.

*Remarks 4.1.* (1) Since we are assuming  $A_i = [0, 1] \times [0, 1] \subset N_i$ , we always have  $m_i < 1$ ,  $i = 1, 2$  (else  $N_i$  would contain a limit cycle).

(2)

Observe that (4.1) continues to hold, though generally with different constants  $\tau_1, \tau_2, m_1, m_2$ , if we replace  $A_1, A_2$  by smaller rectangular neighbourhoods  $A_1^*, A_2^*$  (relative to the same choice of linearizing coordinates). Referring to figure 7, suppose we linearly rescale coordinates on  $A^*$  so that, in the rescaled coordinates,  $A^* = [0, 1] \times [0, 1]$  (we omit the subscript  $i$ ). We may choose unique  $T, S \geq 0$  such that  $\alpha = e^{-\mu S}$ ,  $\beta = e^{-\lambda T}$ . In coordinates  $(x^*, y^*)$  on  $A^* = [0, 1] \times [0, 1]$ , the connection map  $C^* : V^* \rightarrow U^*$  is given by

$$(4.2) \quad C^*(x^*) = e^{(T+S)\lambda} C(e^{-\mu(T+S)} x^*)$$

In particular, if  $C(x) = mx$ , then in rescaled coordinates, we have

$$C^*(x^*) = e^{(T+S)(\lambda-\mu)} m x^*.$$



Since  $\rho_1^j \geq 1$ ,  $j \geq 0$ , the result follows.  $\square$

**Definition 4.4.** Given  $n, m \in \mathbb{N}$  and  $y \in [a, b]$ , we define the closed subinterval  $[Z_m^1(y, n), Z_m^2(y, n)]$  of  $[c, d]$  by

$$[Z_m^1(y, n), Z_m^2(y, n)] = \{z \in [c, d] \mid (y, z) \in E_{n,m}\}.$$

We refer to  $[Z_m^1(y, n), Z_m^2(y, n)]$  as a *bad subinterval*.

**Lemma 4.5.** *If  $n, m \in \mathbb{N}$ ,  $y \in [a, b]$  and  $[Z_m^1(y, n), Z_m^2(y, n)] \neq \emptyset$ , then  $Z_m^1(y, n)$  is given as the maximum of  $c$  and the unique solution of  $s_2^z(m) = s_1^y(n) + \tau_1$  and  $Z_m^2(y, n)$  as the minimum of  $d$  and the unique solution of  $s_2^z(m) = s_1^y(n) - \tau_2$ .*

*Proof.* The result follows by noting that  $s_2^z(m)$  is a decreasing function of  $z \in [c, d]$  and that the endpoints of the bad interval are given as the intersection of  $[c, d]$  with the interval with endpoints determined by the equations  $s_2^z(m) = s_1^y(n) - \tau_2, s_1^y(n) + \tau_1$ .  $\square$

**Proposition 4.6.** *Every bad subinterval  $[Z_m^1(y, n), Z_m^2(y, n)]$  of  $[c, d]$  is contained in the interval  $[\bar{Z}_m^1(y, n), \bar{Z}_m^2(y, n)]$  where,*

$$\begin{aligned} \bar{Z}_m^2(y, n) &= e^{\frac{\lambda_2}{\beta_m}(m\tau_2 - (n-1)\tau_1)} y^{\frac{\lambda_2\alpha_n}{\lambda_1\beta_m}} m_1^{\frac{\lambda_2\pi n}{\lambda_1\theta_m}} m_2^{-\frac{\theta_m}{\beta_m}} \\ \bar{Z}_m^1(y, n) &= e^{\frac{\lambda_2}{\beta_m}((m-1)\tau_2 - n\tau_1)} y^{\frac{\lambda_2\alpha_n}{\lambda_1\beta_m}} m_1^{\frac{\lambda_2\pi n}{\lambda_1\theta_m}} m_2^{-\frac{\theta_m}{\beta_m}} < \bar{Z}_m^2(y, n) \end{aligned}$$

*Proof.* By lemma 4.5,  $[Z_m^1(y, n), Z_m^2(y, n)]$  is contained in an interval with end points given by the solution of  $s_2^z(m) = s_1^y(n) + \tau_1, s_1^y(n) - \tau_2$ . Solving for  $z$ , using lemma 4.2, gives the result.  $\square$

*Remark 4.7.* We have  $\bar{Z}_m^1(y, n) = e^{\frac{-\lambda_2}{\beta_m}(\tau_1 + \tau_2)} \bar{Z}_m^2(y, n)$ .

Let  $y \in [a, b]$ ,  $n \in \mathbb{N}$  and  $t \in [s_1^y(n), s_1^y(n) + 1]$ . The number of turns of  $\phi_t^2(z)$  is an increasing function of  $z \in [c, d]$ . In particular, the maximum number of turns is made when  $z = d$ . If we let  $M_E(n, y)$  denote the number of turns taken by  $d$  around  $\Sigma_2$  in time  $s_1^y(n) + 1$ , we have

$$(4.3) \quad M_E(n, y) = \min\{m \mid s_2^d(m+1) > s_1^y(n) + 1\},$$

Define  $M_E(n) = \sup_{y \in [a, b]} M(n, y)$ .

**Lemma 4.8.** *(Notation and assumptions as above.) If  $n \in \mathbb{N}$ , we have*

- (1)  $M_E(n) < \infty$ .
- (2) *If  $t > 0$  is  $n$ -singular for  $A_1$ , then the number of turns of  $\phi_t^2(z)$  about  $\Sigma_2$  is at most  $M_E(n)$ .*

4.1. **The  $\ell_2$  measure of  $E_\infty$ .** We start by estimating  $\ell_2(E_n)$ .

**Lemma 4.9.** *There exists  $C > 0$  such that for all  $n \in \mathbb{N}$ ,*

$$\ell_2(E_n) \leq C \sum_{m=1}^{\infty} \frac{1}{\lambda_1 \beta_m + \lambda_2 \alpha_n} e^{\frac{\lambda_2}{\beta_m}(\tau_2 m - \tau_1(n-1))}.$$

*Proof.* Let  $n \in \mathbb{N}$ . We have

$$\begin{aligned} \ell_2(E_n) &\leq \int_a^b \sum_{m=1}^{M_E(n)} (\bar{Z}_m^2(y, n) - \bar{Z}_m^1(y, n)) dy \\ &= \int_a^b \sum_{m=1}^{M_E(n)} e^{\frac{\lambda_2}{\beta_m}(\tau_2 m - \tau_1(n-1))} (1 - e^{\frac{-\lambda_2}{\beta_m}(\tau_1 + \tau_2)}) y^{\frac{\lambda_2 \alpha_n}{\lambda_1 \beta_m}} m_1^{\frac{\lambda_2 \pi_n}{\lambda_1 \theta_m}} m_2^{-\frac{\theta_m}{\beta_m}} dy, \\ &= \sum_{m=1}^{M_E(n)} e^{\frac{\lambda_2}{\beta_m}(\tau_2 m - \tau_1(n-1))} (1 - e^{\frac{-\lambda_2}{\beta_m}(\tau_1 + \tau_2)}) m_1^{\frac{\lambda_2 \pi_n}{\lambda_1 \theta_m}} m_2^{-\frac{\theta_m}{\beta_m}} \left( \int_a^b y^{\frac{\lambda_2 \alpha_n}{\lambda_1 \beta_m}} dy \right). \end{aligned}$$

By lemma 4.3, there exists  $C_1 = C_1(m_2) > 0$  such that  $m_2^{-\frac{\theta_m}{\beta_m}} \leq C_1$ , for all  $m \in \mathbb{N}$ . Since  $m_1 < 1$ , we have  $m_1^{\frac{\lambda_2 \pi_n}{\lambda_1 \theta_m}} \leq 1$ , for all  $m, n \in \mathbb{N}$ . We also have

$$\int_a^b y^{\frac{\lambda_2 \alpha_n}{\lambda_1 \beta_m}} dy \leq \int_0^1 y^{\frac{\lambda_2 \alpha_n}{\lambda_1 \beta_m}} dy = \frac{\lambda_1 \beta_m}{\lambda_1 \beta_m + \lambda_2 \alpha_n}.$$

Hence

$$\begin{aligned} \ell_2(E_n) &\leq C_1 \sum_{m=1}^{M_E(n)} \frac{\lambda_1 \beta_m}{\lambda_1 \beta_m + \lambda_2 \alpha_n} e^{\frac{\lambda_2}{\beta_m}(\tau_2 m - \tau_1(n-1))} (1 - e^{\frac{-\lambda_2}{\beta_m}(\tau_1 + \tau_2)}) \\ &\leq C_1 \sum_{m=1}^{M_E(n)} \frac{\lambda_1 \lambda_2 (\tau_1 + \tau_2)}{\lambda_1 \beta_m + \lambda_2 \alpha_n} e^{\frac{\lambda_2}{\beta_m}(\tau_2 m - \tau_1(n-1))}, \end{aligned}$$

where we have used  $(1 - e^{\frac{-\lambda_2}{\beta_m}(\tau_1 + \tau_2)}) \leq (\tau_1 + \tau_2) \frac{\lambda_2}{\beta_m}$ . Hence we have the estimate

$$\ell_2(E_n) \leq C \sum_{m=1}^{\infty} \frac{1}{\lambda_1 \beta_m + \lambda_2 \alpha_n} e^{\frac{\lambda_2}{\beta_m}(\tau_2 m - \tau_1(n-1))}.$$

where  $C = \lambda_1 \lambda_2 C_1 (\tau_1 + \tau_2)$ . □

**Lemma 4.10.**  $\ell_2(E_\infty) = 0$ .

*Proof.* We start by proving that  $\sum_{n=1}^{\infty} \ell_2(E_n) < \infty$ . By lemma 4.9, it suffices to prove that  $\sum_{n,m=1}^{\infty} \frac{1}{\lambda_1\beta_m + \lambda_2\alpha_n} e^{\frac{\lambda_2}{\beta_m}(\tau_2 m - \tau_1(n-1))}$  converges. By the arithmetic-geometric mean inequality we have the estimate

$$\frac{1}{\lambda_1\beta_m + \lambda_2\alpha_n} \leq 1/(2\sqrt{\lambda_1\lambda_2}\sqrt{\beta_m\alpha_n}), \quad m, n \geq 1.$$

It follows easily from the definition of  $\alpha_n, \beta_m$  that there exists  $C > 0$  such that

$$(4.4) \quad \frac{1}{\lambda_1\beta_m + \lambda_2\alpha_n} \leq C\rho_1^{-n/2}\rho_2^{-m/2}, \quad m, n \geq 1.$$

Observe that if  $\tau_2 m - \tau_1(n-1) \leq 0$ , then  $e^{\frac{\lambda_2}{\beta_m}(\tau_2 m - \tau_1(n-1))} \leq 1$ . On the other hand if  $\tau_2 m - \tau_1(n-1) > 0$ , then  $(\tau_2 m - \tau_1(n-1))/\beta_m \leq \tau_2$ , since  $\beta_m \geq m$ , and so  $e^{\frac{\lambda_2}{\beta_m}(\tau_2 m - \tau_1(n-1))} \leq e^{\tau_2\lambda_2}$ . Hence we have a uniform bound on  $e^{\frac{\lambda_2}{\beta_m}(\tau_2 m - \tau_1(n-1))}$  and the convergence of the double sum follows from (4.4). Since  $\sum_{n=1}^{\infty} \ell_2(E_n) < \infty$ , it follows by the Borel-Cantelli lemma [15, Theorem 13.1], that  $\ell_2(E_\infty) = 0$ .  $\square$

**Theorem 4.11.** *The likely limit set of the product of two planar homoclinic attractors  $(\Sigma_1, \phi_t^1)$  and  $(\Sigma_2, \phi_t^2)$  is  $\Sigma$ .*

*Proof.* Our argument for the vanishing of  $\ell_2(E_\infty)$  did not depend on our choices of  $b, d \in [0, 1]$ . We now follow the arguments given at the end of section 3: by  $\sigma$ -additivity of  $\ell_2$ , the set of bad points in  $(\{1\} \times [0, 1]) \times (\{1\} \times [0, 1])$  has measure zero. Using the rescaling argument given in section 3, we deduce that the set of bad points in  $(\{x\} \times [0, 1]) \times (\{x'\} \times [0, 1])$  has measure zero for all  $x, x' \in (0, 1]$ . It follows by Fubini's theorem that the set of bad points in  $A_1 \times A_2$  has measure zero. Hence  $\mathcal{L}(A_1 \times A_2) = \mathcal{L}(N_1 \times N_2) \neq \Sigma_1 \times \Sigma_2$ . and the result follows by theorem 2.6.  $\square$

*Remark 4.12.* Neither of the cycles in the network  $\Sigma \subset \mathbb{R}^4$  is essentially asymptotically stable: for all points (not lying on the stable manifolds), the associated trajectory makes infinitely many traversals arbitrarily close to both homoclinic cycles, even when  $\rho_1 = \rho_2$ .

**4.2. Switching.** In the remainder of the section, we briefly indicate why we cannot expect switching in the network  $\Sigma$ .

Following the notation of Homburg *et al.* [13], let  $\gamma_1 = \Sigma_1 \times \{(0, 0)\}$  and  $\gamma_2 = \{(0, 0)\} \times \Sigma_2$  denote the homoclinic orbits which define the heteroclinic network  $\Sigma$ . Let  $U_\Sigma$  be a tubular neighbourhood of  $\Sigma$  and  $U_0 \subset U_\Sigma$  be a small neighbourhood of  $\mathbf{0}$ . Let  $S_1$  and  $S_2$  be two mutually disjoint cross sections transverse to  $\gamma_1$  and  $\gamma_2$  respectively. We suppose  $S_1, S_2 \subset U_\Sigma \setminus U_0$ .

Let  $\kappa = (k_i) \in \{1, 2\}^{\mathbb{N}}$  be a symbolic sequence. We say that the trajectory with initial condition  $\bar{x}$  is a forward realization of  $\kappa$  if the forward trajectory of  $\bar{x}$  is contained in  $U_\Sigma$  and there exists an increasing sequence of times  $(t_i)_{i \in \mathbb{Z}^+}$ , with  $t_0 = 0$ , such that:

- $\phi_{t_i}^1 \times \phi_{t_i}^2(\bar{x}) \in S_{k_i}, i \in \mathbb{N}$ ;
- $\phi_{t_i}^1 \times \phi_{t_i}^2(\bar{x}) \notin S_1 \cup S_2$ , all  $t \in (t_i, t_{i+1}), i \in \mathbb{N}$ ;
- for  $t \in (t_i, t_{i+1})$ , the trajectory visits  $U_0$  exactly once,  $i \in \mathbb{N}$ .

In other words, a realization of  $\kappa$  is a trajectory that, after an initial transient, follows the homoclinic connections  $\gamma_{k_i}$  in the order prescribed by  $\kappa$ . Our next definition is based on that of Aguiar *et al.* [2] adapted to our context.

**Definition 4.13.** The product of two homoclinic cycles is *switching* if for each symbolic sequence  $\kappa \in \{1, 2\}^{\mathbb{N}}$ , there exists a forward realization of  $\kappa$  in  $U_\Sigma$ .

*Remarks 4.14.* (1) The product of two homoclinic cycles is *finite switching* if the previous set-up holds for finite sequences  $\kappa$  (instead of infinite sequences).

(2) In the obvious way, switching can be defined for the product of two heteroclinic attractors.

**Proposition 4.15.** *The network in  $\Sigma \subset \mathbb{R}^4$  is not switching or finite switching.*

*Proof.* It suffices to prove that  $\Sigma$  is not finite switching. Without loss of generality, suppose that  $U_0 \subset A_1 \times A_2$  and  $K \subset U_0$  is compact and disjoint from  $\Sigma$ . Given  $\bar{x} = (y, z) \in U_0$ , suppose that  $t > 0$  is  $n$ -singular for  $A_1$  (that is, the  $\phi_t^1$  trajectory through  $y$ ). The number of turns of  $\phi_t^2(z)$  about  $\Sigma_2$  is at most  $M_K(n) < \infty$ , where we may choose  $M_K(n)$  independent of  $\bar{x} \in K$ , as in lemma 4.8. All we have to do now is choose a finite sequence where the proportion of 2's grows at a rate faster than  $M(n)/n$ . For example, if we assume  $\rho_1 \leq \rho_2$ , then it is easy to show that there exists  $N \in \mathbb{N}$ , such that  $M_K(n) < 2n$ , for  $n \geq N$ . It follows that if we define the finite block  $\kappa$  by concatenating (1222) sufficiently many times, then  $\kappa$  has no realization.  $\square$ .

## 5. PRODUCTS OF HETEROCLINIC ATTRACTORS

In this section, we study the product of two heteroclinic attractors, both contained in a compact surface. These attractors may be homoclinic cycles, *figure eight* homoclinic cycles, heteroclinic cycles or heteroclinic networks. We only provide detailed computations for the cases where one heteroclinic attractor is a homoclinic loop and

the other is either a heteroclinic cycle with two equilibria or a figure eight homoclinic cycle. The general case is proved along very similar lines (though with more notation).

As we did in the previous section, we assume a simplifying condition on the connection maps (which we remove in section 6).

Let  $M_1, M_2$  denote compact surfaces (possibly with boundary). Suppose that  $\Sigma_1 \subset M_1, \Sigma_2 \subset M_2$  are heteroclinic attractors. Denote the set of equilibria of  $\Sigma_i$  by  $\mathcal{E}_i, i = 1, 2$ . Set  $\Sigma = (\Sigma_1 \times \mathcal{E}_2) \cup (\mathcal{E}_1 \times \Sigma_2)$ .

**Theorem 5.1.** *(Notation and assumptions as above.) The likely limit set of  $\mathcal{B}(\Sigma_1 \times \Sigma_2)$  is the heteroclinic network  $\Sigma$ .*

In order to prove this result it suffices to take the product of any pair of heteroclinic cycles  $\Sigma_1^* \subset \Sigma_1, \Sigma_2^* \subset \Sigma_2$  and show that the likely limit set of  $\mathcal{B}(\Sigma_1^* \times \Sigma_2^*)$  is the heteroclinic network  $\Sigma^* = (\Sigma_1^* \times \mathcal{E}_2^*) \cup (\mathcal{E}_1^* \times \Sigma_2^*)$ , where  $\mathcal{E}_i^* \subset \Sigma_i$  is the set of saddle points on  $\Sigma_i, i = 1, 2$ . Note that we allow for the basin of attraction to be an interior or exterior neighbourhood of the cycle and as well as the cycle being a single homoclinic loop or a figure eight homoclinic cycle. We remark that if  $\Sigma^*$  is a subset of the connected surface  $M$ , then  $M \setminus \Sigma^*$  has either two or three connected components. (The complement of the figure eight homoclinic cycle, figure 1(b), has three connected components as do the sub-cycles of the network shown in figure 1(d). If we allow the underlying manifold to have as its boundary the cycle — allowing corners — then the complement can have one component.)

As indicated above, we only give detailed arguments for the case when  $\Sigma_1^*$  is a planar homoclinic loop and  $\Sigma_2^*$  is either a planar heteroclinic cycle with two equilibria or a planar figure eight homoclinic cycle. Our analysis covers all of the issues that arise in the general case.

**5.1. Product of a homoclinic loop and heteroclinic cycle.** Let  $(\Sigma_1, \phi_t^1, D_1, N_1)$  be a planar attracting homoclinic loop and follow all the notational conventions used in the previous two sections. In particular,  $N_1$  will be an interior neighbourhood of  $\Sigma_1$  and  $\Sigma_1$  will have a hyperbolic saddle point  $\{(0, 0)\} \in \mathbb{R}^2$ . We assume that  $(\Sigma_2, \phi_t^2, D_2, N_2)$  is a planar attracting heteroclinic cycle with hyperbolic equilibria  $\mathbf{p}_1, \mathbf{p}_2$  — see figure 8. As we have drawn this,  $N_2$  will be an interior neighbourhood of  $\Sigma_2$ . Denote the connections between  $\mathbf{p}_1$  and  $\mathbf{p}_2$  by  $\Sigma_{21}, \Sigma_{22}$  so that  $\Sigma_2 = \Sigma_{21} \cup \Sigma_{22}$ .

Suppose the hyperbolic saddle at  $\mathbf{p}_j$  has eigenvalues  $-\mu_2^j < 0 < \lambda_2^j$ , and set  $\eta_1 = \mu_2^1/\lambda_2^1, \eta_2 = \mu_2^2/\lambda_2^2$ . We assume  $\rho_2 = \eta_1\eta_2 > 1$  so that  $\Sigma_2$  is attracting. Since  $\mathbf{p}_1, \mathbf{p}_2$  are hyperbolic saddle points for a  $C^2$  planar




FIGURE 8. Notational conventions for heteroclinic attractor  $\Sigma_2$

flow, we may  $C^1$ -linearize the flow  $\phi_t^2$  on box neighbourhoods  $A_2^1$  of  $\mathbf{p}_1$  and  $A_2^2$  of  $\mathbf{p}_2$ . Set  $A_2 = A_2^1 \cup A_2^2$ . We assume coordinates on  $A_2^1, A_2^2$  are chosen so that equilibria correspond to the origin  $(0, 0) \in \mathbb{R}^2$  and the stable manifold at  $(0, 0)$  is tangent to the  $x$ -axis, the unstable manifold to the  $y$ -axis and  $A_2^j = [-1, 1] \times [-1, 1]$ ,  $j = 1, 2$ . Define

$$U_{21} = \{-1\} \times [0, 1], \quad V_{21} = [0, 1] \times \{-1\},$$

$$U_{22} = \{1\} \times [-1, 0], \quad V_{22} = [-1, 0] \times \{1\}.$$

(See figure 8 and note that  $(V_{21} \times U_{22}) \cup (V_{22} \times U_{21}) \subset N_2$ .) We have  $C^1$  time of first return maps  $T_{21} : V_{21} \rightarrow \mathbb{R}$ ,  $T_{22} : V_{22} \rightarrow \mathbb{R}$  and associated  $C^1$  connection maps  $C_{21} : V_{21} \rightarrow U_{21}$ ,  $C_{22} : V_{22} \rightarrow U_{22}$ . For this section we assume there are strictly positive constants  $\tau_{21}, \tau_{22}, m_{21}, m_{22}$  such that

$$(5.5) \quad T_{21}(x) = \tau_{21}, \quad T_{22}(x) = \tau_{22}, \quad C_{21}(x) = m_{21}x, \quad C_{22}(x) = m_{22}x.$$

After linearly rescaling one coordinate direction if necessary, we may always assume that  $m_{21}, m_{22} < 1$ . In particular, the connection maps  $C_{21}, C_{22}$  are defined on all of  $V_{21}, V_{22}$ .

*Remark 5.2.* If we are given a finite set of planar linear flows  $\phi_t^j$ , each with a hyperbolic saddle point at the origin, box neighbourhoods  $A^j \approx [-1, 1] \times [-1, 1]$ ,  $C^1$  connection maps  $C_j : V_j \rightarrow U_j$ , and  $C^1$  time of first return maps  $T_j : V_j \rightarrow \mathbb{R}$ ,  $j = 1, \dots, r$ , where each product  $V_j \times U_j$  determines a quadrant of  $A^j$ , then this data determines a heteroclinic cycle  $\Sigma$  between  $r$  equilibria. In general,  $\Sigma$  will be defined on a surface, possibly non-orientable. We have chosen the particular configuration

shown in figure 8 because it extends naturally to the exterior of a figure eight attracting homoclinic cycle.

Given  $d \in U_{22} \approx [-1, 0]$ , let  $c > d$  denote the first point of intersection of the forward  $\phi_t^2$ -trajectory through  $d$  with  $U_{22}$ . Necessarily  $[d, c] \subset U_{22}$ . Given  $z \in [d, c]$ , we define strictly monotone increasing sequences  $(t_2^{z,2}(n))_{n \geq 0}$ ,  $(s_2^{z,1}(n))_{n \geq 1}$ ,  $(t_2^{z,1}(n))_{n \geq 1}$ ,  $(s_2^{z,2}(n))_{n \geq 1}$  by

- (1)  $t_2^{z,2}(0) = 0$  and for  $n > 0$ ,  $t_2^{z,j}(n)$  is the time to the  $n$ th intersection of the forward  $\phi_t^2$  trajectory through  $z$  with  $U_{2j}$ ,  $j = 1, 2$ .
- (2)  $s_2^{z,j}(n)$  is the time to the  $n$ th intersection of the forward  $\phi_t^2$  trajectory through  $z$  with  $V_{2j}$ ,  $j = 1, 2$ .

We have

$$0 = t_2^{z,2}(0) < s_2^{z,1}(1) < t_2^{z,1}(1) < s_2^{z,2}(1) < t_2^{z,2}(1) < s_2^{z,1}(2) < \dots$$

Set  $z = z^2(0)$  and for  $n \geq 1$ , let  $z^2(n)$  denote the successive points of intersection of the forward trajectory through  $z$  with  $U_{22}$  ( $z^2(n) = \phi_{t_2^{z,2}(n)}(z)$ ). We similarly let  $z^1(n)$  denote the  $n$ th point of intersection of the forward trajectory through  $z$  with  $U_{21}$  ( $z^1(n) = \phi_{t_2^{z,1}(n)}(z)$ ). A straightforward computation shows that

$$\begin{aligned} z^1(n) &= m_{21}^{(1+\dots+\rho_2^{n-1})} m_{22}^{\eta_1(1+\dots+\rho_2^{n-2})} z^{\eta_1 \rho_2^{n-1}}, \\ z^2(n) &= m_{21}^{\eta_2(1+\dots+\rho_2^{n-1})} m_{22}^{(1+\dots+\rho_2^{n-1})} z^{\rho_2^n}. \end{aligned}$$

As we did in the previous section, it is now straightforward to compute the sequences  $(s_2^{z,j}(n))$ ,  $(t_2^{z,j}(n))$ ,  $j = 1, 2$ .

**Lemma 5.3.** *For  $n \geq 1$  we have*

$$t_2^{z,1}(n) = s_2^{z,1}(n) + \tau_{21}.$$

$$t_2^{z,2}(n) = s_2^{z,2}(n) + \tau_{22}.$$

$$t_2^{z,1}(n) = n\tau_{21} + (n-1)\tau_{22} - \frac{1}{\lambda_2^1} \left( \sum_{j=0}^{n-1} \log z^2(j) \right) - \frac{1}{\lambda_2^2} \left( \sum_{j=1}^{n-1} \log z^1(j) \right).$$

$$t_2^{z,2}(n) = n(\tau_{21} + \tau_{22}) - \frac{1}{\lambda_2^1} \left( \sum_{j=0}^{n-1} \log z^2(j) \right) - \frac{1}{\lambda_2^2} \left( \sum_{j=1}^n \log z^1(j) \right).$$

For  $\rho > 1$ ,  $n \geq 0$ , define

$$\begin{aligned} \pi_n(\rho) &= \frac{\rho^n - n\rho + n - 1}{(\rho - 1)^2}, \\ \alpha_n(\rho) &= \frac{\rho^n - 1}{\rho - 1}. \end{aligned}$$

We have the following expressions for the summations in lemma 5.3.

$$\begin{aligned} \sum_{j=0}^{n-1} \log z^2(j) &= \eta_2 \pi_n(\rho_2) \log m_{21} + \pi_n(\rho_2) \log m_{22} + \alpha_n(\rho_2) \log z \\ \sum_{j=1}^{n-1} \log z^1(j) &= \pi_n(\rho_2) \log m_{21} + \eta_1 \pi_{n-1}(\rho_2) \log m_{22} + \eta_1 \alpha_{n-1}(\rho_2) \log z \end{aligned}$$

**Definition 5.4.** We say  $t > 0$  is  $m$ -singular for  $\Sigma_2$  if

$$t \in [s_2^{z,1}(m), t_2^{z,1}(m)] \cup [s_2^{z,2}(m), t_2^{z,2}(m)].$$

Note that if  $t$  is  $m$ -singular for  $\Sigma_2$ , then  $\phi_t^2(z) \notin A_2^\circ$ .

Set  $E = [a, b] \times [d, c] \subset U_1 \times U_{22}$  (the interval  $[a, b]$  is as defined in the previous section). For  $n, m \in \mathbb{N}$ ,  $j = 1, 2$ , define

$$\begin{aligned} E_{m,n}^j &= \{(y, z) \in E \mid \exists t \in [s_1^y(n), t_1^y(n)] \cap [s_2^{z,j}(n), t_2^{z,j}(n)]\}, \\ E_n^j &= \bigcup_{m=1}^{\infty} E_{m,n}^j, \\ E_n &= E_n^1 \cup E_n^2. \end{aligned}$$

A point  $(y, z) \in E_n$  if there exists  $t > 0$  such that  $t$  is  $n$ -singular for  $\Sigma_1$  and there exists  $m \in \mathbb{N}$  such that  $t$  is  $m$ -singular for  $\Sigma_2$ . We refer to the points of  $E_n$  as the  $n$ -bad points of  $E$ .

Fix  $y \in [a, b]$ . For  $m, n \in \mathbb{N}$ ,  $j = 1, 2$ , define (possibly empty) closed subintervals  $[Z_{j,m}^1(y, n), Z_{j,m}^2(y, n)]$  of  $[d, c]$  by

$$[Z_{j,m}^1(y, n), Z_{j,m}^2(y, n)] = \{z \in [d, c] \mid (y, z) \in E_{m,n}^j\}.$$

We refer to  $[Z_{j,m}^1(y, n), Z_{j,m}^2(y, n)]$  as a bad subinterval.

**Proposition 5.5.** Let  $y \in [a, b]$ . Given  $m, n \in \mathbb{N}$  and  $j = 1, 2$ , we have  $[Z_{j,m}^1(y, n), Z_{j,m}^2(y, n)] \subset [\bar{Z}_{j,m}^1(y, n), \bar{Z}_{j,m}^2(y, n)]$ , where

$$\begin{aligned} \bar{Z}_{1,m}^2(y, n) &= e^{\frac{1}{u_m}(-(n-1)\tau_1 + (m-1)(\tau_{21} + \tau_{22}) + \tau_{21})} y^{\frac{\alpha_n}{u_m \lambda_1}} m_1^{\frac{\pi_n}{u_m \lambda_1}} m_{21}^{-\frac{a_m}{u_m}} m_{22}^{-\frac{b_m}{u_m}}, \\ \bar{Z}_{2,m}^2(y, n) &= e^{\frac{1}{v_m}(-(n-1)\tau_1 + m(\tau_{21} + \tau_{22}))} y^{\frac{\alpha_n}{v_m \lambda_1}} m_1^{\frac{\pi_n}{v_m \lambda_1}} m_{21}^{-\frac{c_m}{v_m}} m_{22}^{-\frac{d_m}{v_m}}, \\ \bar{Z}_{1,m}^1(y, n) &= e^{-\frac{1}{u_m}(\tau_1 + \tau_{21})} \bar{Z}_{1,m}^2(y, n), \\ \bar{Z}_{2,m}^1(y, n) &= e^{-\frac{1}{v_m}(\tau_1 + \tau_{22})} \bar{Z}_{2,m}^2(y, n), \end{aligned}$$

and

$$\begin{aligned} u_m &= \frac{\alpha_m(\rho_2)}{\lambda_2^1} + \frac{\eta_1 \alpha_{m-1}(\rho_2)}{\lambda_2^2}, & v_m &= \frac{\alpha_m(\rho_2)}{\lambda_2^1} + \frac{\eta_1 \alpha_m(\rho_2)}{\lambda_2^2}, \\ a_m &= \frac{\eta_2 \pi_m(\rho_2)}{\lambda_2^1} + \frac{\pi_m(\rho_2)}{\lambda_2^2}, & b_m &= \frac{\pi_m(\rho_2)}{\lambda_2^1} + \frac{\eta_1 \pi_{m-1}(\rho_2)}{\lambda_2^2}, \\ c_m &= \frac{\eta_2 \pi_m(\rho_2)}{\lambda_2^1} + \frac{\pi_{m+1}(\rho_2)}{\lambda_2^2}, & d_m &= \frac{\pi_m(\rho_2)}{\lambda_2^1} + \frac{\eta_1 \pi_m(\rho_2)}{\lambda_2^2}. \end{aligned}$$

*Proof.* For  $j = 1, 2$ , the interval  $[\bar{Z}_{j,m}^1(y, n), \bar{Z}_{j,m}^2(y, n)]$  is contained in an interval with end points given by the solution of  $s_2^{z,j}(m) = s_1^y(n) + \tau_1, s_1^y(n) - \tau_{2j}$ . Solving for  $z$ , using lemma 5.3, gives the result.  $\square$

**Lemma 5.6.** *If we define*

$$S_1 = \sum_{m=1}^{\infty} \frac{1}{u_m \lambda_1 + \alpha_n} e^{\frac{1}{u_m}(-(n-1)\tau_1 + (m-1)(\tau_{21} + \tau_{22}) + \tau_{21})},$$

$$S_2 = \sum_{m=1}^{\infty} \frac{1}{v_m \lambda_1 + \alpha_n} e^{\frac{1}{v_m}(-(n-1)\tau_1 + m(\tau_{21} + \tau_{22}))},$$

then there exists  $C > 0$  such that for all  $n \in \mathbb{N}$ ,

$$\ell_2(E_n) \leq C(S_1 + S_2).$$

*Proof.* As in lemma 4.9, we have

$$\ell_2(E_n) = \ell_1(E_n^1) + \ell_2(E_n^2) \leq \sum_{j=1}^2 \int_a^b \sum_{m=1}^{\infty} \left( \bar{Z}_{j,m}^2(y, n) - \bar{Z}_{j,m}^1(y, n) \right).$$

By lemma 4.3, we have  $0 \leq \frac{a_m}{u_m}, \frac{b_m}{v_m} \leq K_1 \frac{\pi_m(\rho_2)}{\alpha_m(\rho_2)} \leq K_1 \frac{1}{\rho_2 - 1}$ ,  $0 \leq \frac{c_m}{v_m}, \frac{d_m}{v_m} \leq K_2 \frac{\pi_m(\rho_2)}{\alpha_m(\rho_2)} \leq K_2 \frac{1}{\rho_2 - 1}$ , for some constants  $K_1, K_2 > 0$ . Therefore, there exists  $C_1 = C_1(m_1, m_{21}, m_{22}) > 0$  such that  $m_1^{\frac{\pi_n}{u_m \lambda_1}} m_{21}^{-\frac{a_m}{u_m}} m_{22}^{-\frac{b_m}{u_m}} \leq C_1$ ,  $m_1^{\frac{\pi_n}{v_m \lambda_1}} m_{21}^{-\frac{c_m}{v_m}} m_{22}^{-\frac{d_m}{v_m}} \leq C_1$ , for all  $m, n \in \mathbb{N}$ . The remainder of the proof follows that of lemma 4.9.  $\square$

**Lemma 5.7.**  $\ell_2(E_\infty) = 0$ .

*Proof.* It is easy to see that  $u_m, v_m \geq \frac{\alpha_m(\rho_2)}{\lambda_2^1}$ . Hence for some constant  $C > 0$

$$\frac{1}{u_m \lambda_1 + \alpha_n}, \frac{1}{v_m \lambda_1 + \alpha_n} \leq \frac{\lambda_2^1}{\alpha_m(\rho_2) \lambda_1 + \alpha_n \lambda_2^1} \leq C \rho_1^{-n/2} \rho_2^{-m/2}$$

Along similar lines to the proof of lemma 4.1, we show that  $\sum_{n=1}^{\infty} \ell_2(E_n) < \infty$ . It follows by the Borel-Cantelli lemma that  $\ell_2(E_\infty) = 0$ .  $\square$

We have the following special case of theorem 5.1.

**Proposition 5.8.** *The likely limit set of the product of the planar homoclinic cycle  $(\Sigma_1, \phi_t^1)$  and the attracting planar heteroclinic cycle  $(\Sigma_2, \phi_t^2)$  is  $\Sigma$ .*

*Proof.* Similar to that of theorem 4.11.  $\square$

**5.2. Product of a homoclinic loop and a figure eight cycle.**

We now consider the case where  $(\Sigma_2, \phi_t^2, D_2, N_2)$  is an attracting figure eight homoclinic cycle. Write  $N_2 = N_2^- \cup N_2^+$  where  $N_2^- = N_{21}^- \cup N_{22}^-$  is an interior neighbourhood of  $\Sigma_2$  in  $\mathbb{R}^2$  and  $N_2^+$  is an exterior neighbourhood of  $\Sigma_2$  in  $\mathbb{R}^2$  (see figure 9). We may write  $\Sigma_2$  as the union of two attracting homoclinic loops  $\Sigma_{21}, \Sigma_{22}$ .




FIGURE 9. Notational conventions for figure eight homoclinic attractor  $\Sigma_2$

It follows from the results of section 3 applied to the products  $\Sigma_1 \times \Sigma_{21}$  and  $\Sigma_1 \times \Sigma_{22}$ , that  $\mathcal{L}(N_1 \times N_2^-) = \Sigma$ . It remains to show that  $\mathcal{L}(N_1 \times N_2^+) = \Sigma$ .

We suppose that the hyperbolic saddle at  $(0, 0)$  has associated eigenvalues  $-\mu_2 < 0 < \lambda_2$ . Set  $\rho_2 = \mu_2/\lambda_2$  and assume  $\rho_2 > 1$  so that  $\Sigma_2$  is an attracting homoclinic cycle. Since  $(0, 0)$  is a hyperbolic saddle point for a  $C^2$  planar flow, we may  $C^1$ -linearize the flow  $\phi_t^2$  on a box neighbourhood  $A_2 \subset N_2$  of  $(0, 0)$ . We assume coordinates are chosen so that the stable manifold at  $(0, 0)$  is tangent to the  $x$ -axis, the unstable manifold to the  $y$ -axis and  $A_2 = [-1, 1] \times [-1, 1]$ . Define

$$U_{21} = \{-1\} \times [0, 1], \quad V_{21} = [0, 1] \times \{-1\},$$

$$U_{22} = \{1\} \times [-1, 0], \quad V_{22} = [-1, 0] \times \{1\}.$$

(See inset to figure 9.)

But now we are exactly in the situation described by figure 8. Our previous results extend immediately (with  $\lambda_2 = \lambda_2^1 = \lambda_2^2$ ,  $\mu_2 = \mu_2^1 = \mu_2^2$  and the terms  $\eta_1, \eta_2$  replaced by  $\rho_2 = \mu_2/\lambda_2$ , and  $\rho_2$  by  $\rho_2^2$ ,  $m_2 = m_{21} = m_{22}$ ,  $\tau_2 = \tau_{21} = \tau_{22}$ ).

Summarizing, we have shown

**Proposition 5.9.** *The likely limit set of the product of a planar homoclinic attractor  $(\Sigma_1, \phi_t^1)$  and an attracting planar figure eight cycle  $(\Sigma_2, \phi_t^2)$ , is  $\Sigma$ .*

## 6. GENERAL GLOBAL MAPS

In the previous sections, we assumed that the connection maps  $C : V \rightarrow U$  were linear and the associated time maps  $T : V \rightarrow \mathbb{R}$  were constant. In this section, we remove this restriction and show that our results continue to hold. We give the details only for product of two attracting homoclinic loops (details for the general case of products of heteroclinic attractors are similar and use the same methods). As far as possible we follow the notational conventions of sections 3, 4. The reader should note that a particular concern is to obtain results where we can use the rescaling strategy described at the end of section 3.

Since we are assuming flows are at least  $C^2$ , the maps  $C_i : V_i \rightarrow U_i$ ,  $T_i : V_i \rightarrow \mathbb{R}$  defined in section 3 are  $C^2$  (see Wiggins [24, Section 10.3]). Hence for  $i = 1, 2$  we may write

$$(6.6) \quad C_i(x) = \gamma^i(x)x$$

$$(6.7) \quad T_i(x) = \tau_i + O(|x|),$$

where  $\gamma^i$  is  $C^1$ ,  $m_i = \gamma^i(0) \in (0, 1)$  and  $\tau_i = T_i(0)$ .

Variation in the time maps  $T_i$  causes only minor problems. However, the case of general connection maps is delicate — this is already evident in the analysis given in [4, §3] for the product of a homoclinic loop and limit cycle. We start with an analysis of the case when  $\rho_1 \neq \rho_2$  and conclude with the harder case  $\rho_1 = \rho_2$ . Note that the results in the first part of the section hold with no restriction on  $\rho_1, \rho_2 (> 1)$ . We indicate in the text results which do not hold with  $\rho_1 = \rho_2$ .

*Rescaling.* We start with an analysis of how the connection map changes when we restrict to a smaller rectangular  $\hat{A} \subset A$  (we omit subscripts in what follows). Let  $A = [0, 1] \times [0, 1]$  and  $\hat{A} = [0, \alpha] \times [0, \beta] \subset A$ . Let  $\alpha = e^{-\mu S}$ ,  $\beta = e^{-\lambda T}$ . Linearly rescale coordinates on  $A$  so that, in the new coordinates  $(\hat{x}, \hat{y})$ ,  $\hat{A} = [0, 1] \times [0, 1]$ . If we let  $\hat{C} : V^* \rightarrow U^*$  denote the connection map associated to  $\hat{A}$ , we have

$$\hat{C}(\hat{x}) = e^{(T+S)\lambda} C(e^{-\mu(T+S)} \hat{x})$$

(See remarks 4.1(2).) Writing  $C(x) = \gamma(x)x$ , gives

$$(6.8) \quad \hat{\gamma}(\hat{x}) = e^{(T+S)(\lambda-\mu)}\gamma(e^{-(T+S)\mu}\hat{x})$$

$$(6.9) \quad \frac{\hat{\gamma}'(\hat{x})}{\hat{\gamma}(\hat{x})} = e^{-(T+S)\mu} \frac{\gamma'(e^{-(T+S)\mu}\hat{x})}{\gamma(e^{-(T+S)\mu}\hat{x})}$$

Let  $c_1 = \sup_{x \in [0,1]} |\gamma'(x)|$ .

**Lemma 6.1.** *Let  $T, S \geq 0$  and define  $\hat{m} = \hat{\gamma}(0)$ ,  $\hat{m}^+ = \sup_{\hat{x} \in [0,1]} \hat{\gamma}(\hat{x})$ ,  $\hat{m}^- = \inf_{\hat{x} \in [0,1]} \hat{\gamma}(\hat{x})$ . We have*

$$(1) \quad \hat{m} = e^{(T+S)(\lambda-\mu)}m.$$

$$(2) \quad \hat{m}(1 - \frac{c_1}{m}e^{-\mu(T+S)}) \leq \hat{m}^- \leq \hat{m}^+ \leq \hat{m}(1 + \frac{c_1}{m}e^{-\mu(T+S)}).$$

*These estimates continue to hold if we increase either  $T$  or  $S$ .*

*Proof.* The estimates follow straightforwardly from (6.8,6.9).  $\square$

Define  $\varepsilon_0 > 0$  by

$$\varepsilon_0 = \min\{1/4, (\rho_2 - 1)/4\}.$$

It follows from lemma 6.1 and (6.9) that we can choose  $A_2$  so that

$$(6.10) \quad m_2^+ \leq \min\{\rho_2^{-2/\rho_2}, 2^{-3}\}.$$

$$(6.11) \quad \left| \frac{\gamma'(x)}{\gamma(x)} \right| \leq \frac{\varepsilon_0}{2}.$$

Shrinking  $A_i$  if necessary, we can assume that there exist  $\tau_i^-, \tau_i^+ > 0$  so that  $\tau_i^- \leq T_i(x) \leq \tau_i^+$ , for all  $x \in V_i$ .

*Remarks 6.2.* (1) If we shrink  $A_2$  further then (6.10,6.11) continue to hold —  $m_2^+$  is a decreasing function of  $T + S$ .

(2) If we shrink  $A_i$ ,  $\tau_i^\pm$  increase,  $i = 1, 2$ .

*Defining the sequences.* Exactly as in section 3, choose a rectangle  $E = [a, b] \times [c, d] \subset U_1 \times U_2$ . Given  $y = y_0 \in U_1$ ,  $z = z_0 \in U_2$  let  $(y_n)$  and  $(z_n)$  denote the successive points of intersection of the forward trajectories through  $y_0$  with  $U_1$  and through  $z_0$  with  $U_2$ . For  $n \geq 1$ , we have

$$y_n = \gamma^1(y_{n-1}^{\rho_1})y_{n-1}^{\rho_1}, \quad z_n = \gamma^2(z_{n-1}^{\rho_2})z_{n-1}^{\rho_2}.$$

For  $n \geq 0$  define  $\gamma_n^1(y) = \gamma^1(y_n)$ ,  $\gamma_n^2(z) = \gamma^2(z_n)$ .

We have a straightforward generalization of lemma 4.2.

**Lemma 6.3.** *Let  $(y, z) \in E$ ,  $n \in \mathbb{N}$ . We have*

$$(1) \quad t_1^y(n) = s_1^y(n) + T_1(y_n^{\rho_1}) \text{ and } t_2^z(n) = s_2^z(n) + T_2(z_n^{\rho_2}).$$

$$(2) \quad t_1^y(n) = \sum_{i=0}^{n-1} T_1(y_i^{\rho_1}) - \frac{1}{\lambda_1}(\alpha_n \log y + \log \prod_{j=0}^{n-2} (\gamma_j^1(y))^{\alpha_{n-1-j}}).$$

$$(3) \quad t_2^z(n) = \sum_{i=0}^{n-1} T_2(z_i^{\rho_2}) - \frac{1}{\lambda_2}(\beta_n \log z + \log \prod_{j=0}^{n-2} (\gamma_j^2(z))^{\beta_{n-1-j}}).$$

*Remark 6.4.* If we assume the connection maps are linear,  $\gamma^i = m_i$ ,  $i = 1, 2$ , we find that

$$\prod_{j=0}^{n-2} (\gamma_j^1(y))^{\alpha_{n-1-j}} = m_1^{\sum_{j=0}^{n-2} \alpha_{n-1-j}} = m_1^{\pi_n},$$

with a similar expression for  $\prod_{j=0}^{n-2} (\gamma_j^2(z))^{\beta_{n-1-j}}$ . These are the terms that appear in lemma 4.2.

*The main estimate.* In this section we derive the main estimate on  $|(\gamma_n^2)'(z)/\gamma_n^2(z)|$  that we use for our proof of convergence of  $\sum \ell_2(E_n)$ . So as to simplify the notation, we generally drop the identifying subscript 2 from  $\gamma_j^2, \gamma_2$  and  $\rho_2$ . Given  $n \in \mathbb{N}$ , define  $\bar{\gamma}_n : [c, d] \rightarrow \mathbb{R}$  by

$$z_n = \bar{\gamma}_n(z)z^{\rho^n}, \quad z \in [c, d].$$

Since we have  $\gamma(x)x \leq m^+x$ , for all  $x \in (0, 1]$ , we have the easy estimate

$$(6.12) \quad z_n \leq (m^+)^{\beta_n} z^{\rho^n}, \quad n \geq 1,$$

where  $\beta_n = \sum_{j=0}^{n-1} \rho^j$ .

**Lemma 6.5.** *For  $n \geq 1$  we have*

- (1)  $\left| \frac{\bar{\gamma}'_n(z)}{\bar{\gamma}_n(z)} \right| \leq \varepsilon_0 \rho^n.$
- (2)  $\left| \frac{\gamma'_n(z)}{\gamma_n(z)} \right| \leq \varepsilon_0.$

If  $S \geq 0$  and we linearly rescale coordinates in the  $x$ -direction so that  $e^{-\mu S}$  is rescaled to 1, then estimate (2) changes to

$$\left| \frac{\gamma'_n(z)}{\gamma_n(z)} \right| \leq e^{-\mu S} \varepsilon_0.$$

*Proof.* The proof of (1) is very similar to that of lemma 3.4 in [4] and we only indicate the main points. The proof goes by induction on  $n$  with the hypothesis at step  $n$  being

$$\left| \frac{\bar{\gamma}'_n(z)}{\bar{\gamma}_n(z)} \right| \leq \varepsilon_0(1 - 2^{-n})\rho^n.$$

When  $n = 1$ , we have  $\bar{\gamma}_1(z) = \gamma(z^\rho)$  and differentiating gives

$$\frac{\bar{\gamma}'_1(z)}{\bar{\gamma}_1(z)} = \rho z^{\rho-1} \frac{\gamma'(z^\rho)}{\gamma(z^\rho)}.$$

Hence by (6.11), we have  $|\rho z^{\rho-1} \frac{\gamma'(z^\rho)}{\gamma(z^\rho)}| \leq \varepsilon_0 \frac{\rho}{2} z^{\rho-1} \leq \varepsilon_0 \frac{\rho}{2}$ , since  $z \in (0, 1]$ . This verifies the result when  $n = 1$ . Given the truth of the statement for  $n - 1$ , the proof of step  $n$  proceeds by estimating the derivative  $\bar{\gamma}'_n$  and uses the bound  $m_2^+ \leq 2^{-3}$  (see (6.10)) and (6.12).

Turning to (2), we have  $z_n = \bar{\gamma}_n(z)z^{\rho^n}$  and differentiating with respect to  $z$  gives

$$z'_n = \bar{\gamma}'_n(z)z^{\rho^n} + \bar{\gamma}_n(z)\rho^n z^{\rho^n-1}.$$

We have  $\gamma_n(z) = \gamma(z_n^\rho)$ . Differentiating and dividing by  $\gamma_n(z)$ ,

$$\begin{aligned} \frac{\gamma'_n(z)}{\gamma_n(z)} &= \frac{\gamma'(z_n^\rho)}{\gamma(z_n^\rho)} \rho z_n^{\rho-1} z'_n, \\ &= \frac{\gamma'(z_n^\rho)}{\gamma(z_n^\rho)} \rho z_n^{\rho-1} (\bar{\gamma}'_n(z)z^{\rho^n} + \bar{\gamma}_n(z)\rho^n z^{\rho^n-1}). \end{aligned}$$

By (6.11), we have

$$\left| \frac{\gamma'_n(z)}{\gamma_n(z)} \right| \leq \varepsilon_0 \frac{\rho z_n^{\rho-1}}{2} (|\bar{\gamma}'_n(z)z^{\rho^n}| + |\bar{\gamma}_n(z)\rho^n z^{\rho^n-1}|).$$

By (1) and choice of  $\varepsilon_0$ , we have  $|\bar{\gamma}'_n(z)| \leq \varepsilon_0 \rho^n \bar{\gamma}_n(z) \leq \rho^n \bar{\gamma}_n(z)$  and so

$$\begin{aligned} \frac{\rho z_n^{\rho-1}}{2} |\bar{\gamma}'_n(z)z^{\rho^n}| &\leq \frac{\rho z_n^{\rho-1}}{2} \rho^n \bar{\gamma}_n(z) z^{\rho^n}, \\ &= \frac{\rho^{n+1} z_n^\rho}{2}, \\ &\leq \frac{\rho^{n+1} (m^+)^{\rho\beta_n}}{2}. \end{aligned}$$

Turning to the second term, we have

$$\begin{aligned} \frac{\rho z_n^{\rho-1}}{2} |\bar{\gamma}_n(z)\rho^n z^{\rho^n-1}| &= \frac{z_n^\rho \rho^{n+1}}{2z}, \\ &\leq \frac{(m^+)^{\rho\beta_n} z^{\rho^{n+1}} \rho^{n+1}}{2z}, \\ &\leq \frac{\rho^{n+1} (m^+)^{\rho\beta_n}}{2}. \end{aligned}$$

Using the bound  $m^+ \leq \rho^{-2/\rho}$  (see (6.10)), we get  $\rho^{n+1} (m^+)^{\rho\beta_n} \leq 1$ .

The final statement is immediate from (6.9).  $\square$

*Remarks 6.6.* (1) Lemma 6.5 continues to hold, with the smaller constants, if we shrink  $A_2$ .

(2) Estimate (2) of lemma 6.10 will more than suffice to handle the situation when  $\rho_1 \neq \rho_2$ . The final statement will be crucial for the analysis when  $\rho_1 = \rho_2$ .

*Estimating  $\ell_2(E_n)$  when  $\rho_2 > \rho_1$ .* For this section we assume  $\rho_2 \neq \rho_1$ . Without loss of generality we suppose  $\rho_2 > \rho_1$ . Under these conditions, we prove that there exists  $N \in \mathbb{N}$ ,  $C > 0$  such that for all  $n \geq N$ ,

$$(6.13) \quad \ell_2(E_n) \leq C \frac{n}{\alpha_n}$$

Since  $\sum_{n \geq 1} \frac{n}{\alpha_n} < \infty$ , this implies that  $\sum_{n \geq 1} \ell_2(E_n) < \infty$  and so  $\ell_2(E_\infty) = 0$ .

We continue to assume  $(y, z) \in E = [a, b] \times [c, d]$ . Define

$$M_1 = M_1(n, y) = \prod_{j=0}^{n-2} (\gamma_j^1(y))^{\alpha_{n-1-j}}, \quad M_2(n, z) = \prod_{j=0}^{n-2} (\gamma_j^2(z))^{\beta_{n-1-j}}$$

$$N_1 = N_1(n, y) = \sum_{i=0}^{n-2} T_1(y_i^{\rho_1}), \quad N_2(n, z) = \sum_{i=0}^{n-2} T_2(z_i^{\rho_2})$$

The bad intervals are contained in intervals  $[Z_m^1(y, n), Z_m^2(y, n)]$  which have nonempty intersection with  $[c, d]$ . The points of  $Z_m^1(y, n), Z_m^2(y, n)$  are obtained by solving the following equations for  $z$

$$(6.14) \quad s_1^y(n) = s_2^z(m) + T_2(z_m^{\rho_2}), \quad s_1^y(n) = s_2^z(m) - T_1(y_n^{\rho_1}).$$

For  $i = 1, 2$  define

$$(6.15) \quad N_{2i} = N_2(m, Z_m^i(y, n)), \quad M_{2i} = M_2(m, Z_m^i(y, n))$$

Using (6.14), we find that

$$Z_m^1(y, n) = e^{\frac{\lambda_2}{\beta_m}(N_{21} - N_1 + T_2(z_m^{\rho_2}))} y^{\frac{\lambda_2 \alpha_n}{\lambda_1 \beta_m}} M_1^{\frac{\lambda_2}{\lambda_1 \beta_m}} M_{21}^{-\frac{1}{\beta_m}}$$

$$Z_m^2(y, n) = e^{\frac{\lambda_2}{\beta_m}(N_{22} - N_1 - T_1(y_n^{\rho_1}))} y^{\frac{\lambda_2 \alpha_n}{\lambda_1 \beta_m}} M_1^{\frac{\lambda_2}{\lambda_1 \beta_m}} M_{22}^{-\frac{1}{\beta_m}}$$

The maximum number of turns around  $\Sigma_2$  in time  $s_1^y(n) + \tau_1^+$  is taken when  $z = d$ , and is estimated by

$$(6.16) \quad M_E(n, y) = \min\{m \mid s_2^d(m+1) > s_1^y(n) + \tau_1^+\}$$

Define  $M_E(n) = \sup_{y \in [a, b]} M_E(n, y)$ . Choose  $r > 0$  such that  $\rho_2^r < \rho_1$ .

**Lemma 6.7.** *There exists  $N_1 \in \mathbb{N}$  such that  $M_E(n) \leq rn$  for all  $n \geq N_1$ .*

*Proof.* By definition,  $s_2^d(M_E(n) + 1) > s_1^y(n) + \tau_1^+$ . The result now follows by a straightforward computation using the expressions for  $s_2^d(M_E(n) + 1)$  and  $s_1^y(n)$  given by lemma 6.3. Indeed, the condition  $\rho_2 > \rho_1$  implies that  $\lim_{n \rightarrow \infty} M_E(n)/n = 0$ .  $\square$

Let  $J(n) = \{m \in \mathbb{N} \mid [Z_m^1(y, n), Z_m^2(y, n)] \cap [c, d] \neq \emptyset\}$ . For  $n \in \mathbb{N}$ , define

$$K_n = \sup_{m \in J(n)} \left\{ e^{-\frac{\lambda_2}{\beta_m}((n-1)\tau_1^- - m\tau_2^+)} (m_1^+)^{\frac{\pi_n \lambda_2}{\lambda_1 \beta_m}} (m_2^-)^{-\frac{\theta_m}{\beta_m}} \right\}$$

**Lemma 6.8.** *There exists  $N \geq N_1$  such that*

$$(6.17) \quad K_n \leq 2, \quad n \geq N.$$

*Proof.* If  $n \geq N_1$ , then for all  $m \in J(n)$ ,  $m \leq rn$ . It follows easily from the condition  $\rho_2^r < \rho_1$  and lemma 4.3 that  $\lim_{n \rightarrow \infty} K_n = 0$ .  $\square$

*Remark 6.9.* Lemma 6.8 does not extend to the case when  $\rho_2 = \rho_1$ . The difficulty lies with the term  $(m_2^-)^{-\frac{\theta_m}{\beta_m}}$  which we can only bound by  $(m_2^-)^{-\frac{1}{\rho_2^2 - 1}}$  (lemma 4.3). We handle this problem at the end of the section.

For  $n \in \mathbb{N}$ ,  $m \in J(n)$  set  $D_{m,n}^y = Z_m^2(n, y) - Z_m^1(n, y)$ ,  $m \in J(n)$ . We have

$$\ell_2(E_n) \leq \int_a^b \sum_{m=1}^{M_E(n)} D_{m,n}^y dy$$

**Lemma 6.10.** *For  $n \in \mathbb{N}$ ,  $m \in J(n)$*

$$(6.18) \quad D_{m,n}^y \leq (I_1 + I_2 + I_3) y^{\frac{\lambda_2 \alpha_n}{\lambda_1 \beta_m}}$$

where

$$I_1 = K_n \frac{\lambda_2}{\beta_m} (m-1)(\tau_2^- - \tau_2^+), \quad I_2 = K_n \frac{\lambda_2}{\beta_m} (\tau_1^+ + \tau_2^+) \quad I_3 = K_n \left| \log \left( \frac{M_{22}}{M_{21}} \right) \right|.$$

*Proof.* We have

$$\begin{aligned} D_{m,n}^y &\leq e^{-\frac{\lambda_2}{\beta_m} N_1} y^{\frac{\lambda_2 \alpha_n}{\lambda_1 \beta_m}} M_1^{\frac{\lambda_2}{\lambda_1 \beta_m}} e^{\frac{\lambda_2}{\beta_m} ((m-1)\tau_2^+)} e^{\frac{\lambda_2}{\beta_m} \tau_2^+} (M_{21})^{-\frac{1}{\beta_m}} \\ &\quad - e^{-\frac{\lambda_2}{\beta_m} N_1} y^{\frac{\lambda_2 \alpha_n}{\lambda_1 \beta_m}} M_1^{\frac{\lambda_2}{\lambda_1 \beta_m}} e^{\frac{\lambda_2}{\beta_m} ((m-1)\tau_2^-)} e^{-\frac{\lambda_2}{\beta_m} \tau_1^+} (M_{22})^{-\frac{1}{\beta_m}} \\ &\leq e^{-\frac{\lambda_2}{\beta_m} N_1} y^{\frac{\lambda_2 \alpha_n}{\lambda_1 \beta_m}} (m_1^+)^{\frac{\pi_n \lambda_2}{\lambda_1 \beta_m}} e^{\frac{\lambda_2}{\beta_m} ((m-1)\tau_2^+)} e^{\frac{\lambda_2}{\beta_m} \tau_2^+} (m_2^-)^{-\frac{\theta_m}{\beta_m}} \\ &\quad \times \left( 1 - e^{\frac{\lambda_2}{\beta_m} ((m-1)(\tau_2^- - \tau_2^+) - (\tau_1^+ + \tau_2^+))} \left( \frac{M_{22}}{M_{21}} \right)^{-\frac{1}{\beta_m}} \right) \\ &\leq K_n \left( \frac{\lambda_2}{\beta_m} ((m-1)(\tau_2^+ - \tau_2^-) + (\tau_1^+ + \tau_2^+)) + \frac{1}{\beta_m} \log \left( \frac{M_{22}}{M_{21}} \right) \right) y^{\frac{\lambda_2 \alpha_n}{\lambda_1 \beta_m}} \end{aligned}$$

By separating the three terms, we obtain the result.  $\square$

*Remark 6.11.* In the case when  $\tau_2^+ = \tau_2^- = \tau_2$ ,  $M_{21} = M_{22} = m_2^{\theta_m}$ , the terms  $I_1$  and  $I_3$  vanish, and we are only left with  $I_2$  which is precisely the term in lemma 4.9. The crucial term we have to estimate is  $\left| \log \left( \frac{M_{22}}{M_{21}} \right) \right|$  in  $I_3$ .

**Lemma 6.12.** *We have  $\left| \log \left( \frac{M_{22}}{M_{21}} \right) \right| \leq \varepsilon_0 \frac{\beta_m}{\rho_2 - 1} D_{m,n}^y$ .*

*Proof.* Using (6.15), we get

$$\begin{aligned}
\left| \log \left( \frac{M_{22}}{M_{21}} \right) \right| &= \left| \log \prod_{j=0}^{m-2} \left( \frac{\gamma_j(Z_m^2(n, y))}{\gamma_j(Z_m^1(n, y))} \right)^{\beta_{m-j-1}} \right|, \\
&\leq \sum_{j=0}^{m-2} \beta_{m-j-1} \left| \log \frac{\gamma_j(Z_m^2(n, y))}{\gamma_j(Z_m^1(n, y))} \right|, \\
&\leq \sum_{j=0}^{m-2} \beta_{m-j-1} \left| \frac{\gamma_j'(w)}{\gamma_j(w)} \right| D_{m,n}^y, \\
&\leq \varepsilon_0 \sum_{j=0}^{m-2} \beta_{m-j-1} D_{m,n}^y, \quad (\text{lemma 6.5}), \\
&= \varepsilon_0 \frac{1}{\rho_2 - 1} \sum_{j=0}^{m-2} (\rho_2^{m-j-1} - 1) D_{m,n}^y \\
&\leq \varepsilon_0 \frac{1}{\rho_2 - 1} \sum_{j=0}^{m-2} \rho_2^{m-j-1} D_{m,n}^y \\
&\leq \varepsilon_0 \frac{\beta_m}{\rho_2 - 1} D_{m,n}^y.
\end{aligned}$$

The second inequality is obtained by using mean value theorem on the interval  $[Z_m^1(n, y), Z_m^2(n, y)]$ ,  $w \in (Z_m^1(n, y), Z_m^2(n, y))$ .  $\square$

**Lemma 6.13.** *For  $n \geq N$ , there exists  $C > 0$  such that*

$$\ell_2(E_n) \leq C \frac{n}{\alpha_n}$$

*Proof.* We have

$$I_1 \leq C_1 \frac{m}{\beta_m} y^{\frac{\lambda_2 \alpha_n}{\lambda_1 \beta_m}}, \quad I_2 = C_2 \frac{1}{\beta_m} y^{\frac{\lambda_2 \alpha_n}{\lambda_1 \beta_m}}, \quad I_3 \leq C_3 y^{\frac{\lambda_2 \alpha_n}{\lambda_1 \beta_m}} D_{m,n}^y,$$

where  $C_1 = K_n \lambda_2 (\tau_2^+ - \tau_2^-)$ ,  $C_2 = K_n \lambda_2 (\tau_1^+ + \tau_2^+)$ ,  $C_3 = \varepsilon_0 K_n / (\rho_2 - 1)$ .

*Proof.* Since  $D_{m,n}^y \leq I_1 + I_2 + I_3$  we have

$$\begin{aligned} D_{m,n}^y &\leq (mC_1 + C_2) \frac{1}{\beta_m} y^{\frac{\lambda_2 \alpha_n}{\lambda_1 \beta_m}} + C_3 y^{\frac{\lambda_2 \alpha_n}{\lambda_1 \beta_m}} D_{m,n}^y \\ &\leq (C_1 + C_2) \frac{m}{\beta_m} y^{\frac{\lambda_2 \alpha_n}{\lambda_1 \beta_m}} + C_3 y^{\frac{\lambda_2 \alpha_n}{\lambda_1 \beta_m}} D_{m,n}^y \\ &\leq (C_1 + C_2) \frac{m}{\beta_m} y^{\frac{\lambda_2 \alpha_n}{\lambda_1 \beta_m}} + \frac{1}{2} y^{\frac{\lambda_2 \alpha_n}{\lambda_1 \beta_m}} D_{m,n}^y, \end{aligned}$$

where the last inequality holds by lemma 6.8 for  $n \geq N$  and our choice of  $\varepsilon_0 = \min\{1/4, (\rho_2 - 1)/4\}$ . Hence for  $n \geq N$ , we have the estimate

$$\begin{aligned} D_{m,n}^y &\leq (C_1 + C_2) \frac{m}{\beta_m} \frac{y^{\frac{\lambda_2 \alpha_n}{\lambda_1 \beta_m}}}{1 - y^{\frac{\lambda_2 \alpha_n}{\lambda_1 \beta_m} / 2}} \\ &\leq \left( \frac{2m\lambda_1(C_1 + C_2)}{\lambda_2 \alpha_n} \right) \frac{\frac{\lambda_2 \alpha_n}{\lambda_1 \beta_m} y^{\frac{\lambda_2 \alpha_n}{\lambda_1 \beta_m} - 1} / 2}{1 - y^{\frac{\lambda_2 \alpha_n}{\lambda_1 \beta_m} / 2}} \end{aligned}$$

Integrating, we get

$$\begin{aligned} \int_{y=a}^b D_{m,n}^y dy &\leq \int_0^1 D_{m,n}^y dy \\ &= \left( \frac{2m\lambda_1(C_1 + C_2)}{\lambda_2 \alpha_n} \right) \log 2 \\ &\leq C \frac{n}{\alpha_n}. \end{aligned}$$

where  $C = \frac{2r\lambda_1(C_1+C_2)}{\lambda_2}$ . □

*Remark 6.14.* Our arguments extend immediately to the case of rectangles  $E \subset (\{x\} \times [0, 1]) \times (\{x'\} \times [0, 1])$ , where  $x \in V_1, x' \in V_2$ . The key observation is that lemma 6.7 continues to hold and so our arguments all continue to apply with the *same*  $\varepsilon_0$ .

*Estimating  $\ell_2(E_n)$  when  $\rho_2 = \rho_1$ .* Assume that  $\rho_1 = \rho_2 = \rho$ . We need replacements for lemmas 6.7, 6.8. Following our earlier notation, we may choose  $N_1 = N(\rho, \lambda_2, \tau_1^-, \tau^+) \in \mathbb{N}$  such that

$$\sup_{n \geq N_1} K_n \leq 2(m_2^-)^{-\frac{1}{\rho-1}}.$$

Set  $k = m_2^-$ . Linearly rescale  $A_2$  in the  $x$ -direction by  $e^{-\mu_2 S}$ , where  $S$  is chosen so that

$$(6.19) \quad e^{S(\lambda_2 - \mu_2)} k^{-\frac{1}{\rho-1}} \leq 1.$$

If we let  $\hat{A}_2 = [0, e^{-\mu_2 S}] \times [0, 1]$ , then a lower bound for  $\hat{m}_2^-$  on  $\hat{A}_2$  is  $e^{S(\lambda_2 - \mu_2)}k$  and the constant  $\varepsilon_0$  in lemma 6.5 rescales to  $\hat{\varepsilon}_0 = \varepsilon_0 e^{-\mu S}$ .

Replace  $A_2$  by  $\hat{A}_2$ . On the rescaled  $A_2$  we have

$$\begin{aligned} (m_2^-)^{-\frac{1}{\rho-1}} &\leq (k e^{S(\lambda_2 - \mu_2)})^{-\frac{1}{\rho-1}} = k^{-\frac{1}{\rho-1}} e^{S\lambda_2}, \\ \varepsilon_0 &= e^{-\mu S} \min\left\{\frac{1}{4}, \frac{\rho-1}{4}\right\}, \\ \sup_{n \geq N_1} K_n &\leq 2k^{-\frac{1}{\rho-1}} e^{S\lambda_2}. \end{aligned}$$

Lemma 6.12 holds without any assumptions on  $\rho_1, \rho_2$  and so we still have  $\left| \log \left( \frac{M_{22}}{M_{21}} \right) \right| \leq \varepsilon_0 \frac{\beta_m}{\rho-1} D_{m,n}^y$ . We have the following straightforward variation on lemma 6.7.

**Lemma 6.15.** *If  $\rho_1 = \rho_2$ , there exists  $N \geq N_1$  such that*

$$M_E(n) \leq 2n, \quad n \geq N.$$

Finally, for lemma 6.13 to hold, it is enough that  $\varepsilon_0 K_n / (\rho - 1) \leq \frac{1}{2}$ ,  $n \geq N$ . Computing, we find that

$$\begin{aligned} \varepsilon_0 K_n / (\rho - 1) &\leq 2 \min\left\{\frac{1}{4}, \frac{\rho-1}{4}\right\} e^{-\mu_2 S} k^{-\frac{1}{\rho-1}} e^{S\lambda_2} / (\rho_2 - 1) \\ &\leq \frac{2}{\rho-1} \min\left\{\frac{1}{4}, \frac{\rho-1}{4}\right\}, \text{ by (6.19),} \\ &\leq \frac{1}{2}. \end{aligned}$$

*Remark 6.16.* Of course, we could use our argument for the resonant case  $\rho_1 = \rho_2$  in the general case. However, we prefer to present the arguments separately as we feel both arguments have intrinsic interest and may be relevant in problems where product structure is broken.

## 7. SIMULATIONS

In this section, we present numerical simulations which illustrate our results on the likely limit set for the product flow of two attracting homoclinic orbits. We consider the product of identical dynamical systems given by

$$(7.20) \quad x' = y - x + (x - a)^2,$$

$$(7.21) \quad y' = y + 3x - x^3,$$

where  $a = 1.7611050$ . The two-dimensional system has an attracting homoclinic cycle  $\Sigma_1$  associated to the saddle point  $p = (0.452, -1.263)$  (see figure 10). Since the attractors are identical for the product system, we see that if the initial conditions are equal then the  $\omega$ -limit

set will be the diagonal  $\{(u, u), u \in \Sigma_1\} \subset \Sigma_1 \times \Sigma_1$ . We look at the  $\omega$ -limit set of the trajectory with initial condition  $((x_1, y_1), (x_2, y_2)) = ((2, 2.1), (1.7, 1))$ . Ignoring the initial transient, we show the projections of this trajectory on the  $(x_1, x_2)$ - and  $(y_1, y_2)$ -planes in figures 11 and 12.

The projections of the product flow on the  $(x_1, x_2)$ - and  $(y_1, y_2)$ -planes are illustrated in figures 11 and 12. It is clear from figures that when the  $(x_1, y_1)$  components of the trajectory are in the neighbourhood of  $p$  then the  $(x_2, y_2)$  components of the trajectory visits each point of the homoclinic orbit, and vice-a-versa. The results of the simulation are consistent with the  $\omega$ -limit being equal to  $\Sigma$  and not  $\Sigma_1 \times \Sigma_1$ . (Simulations were performed using XPPAUT software [9].)






FIGURE 10. Projection of  $\omega$ -limit set onto  $x_1, y_1$  plane.

## 8. BIFURCATIONS NEAR THE PRODUCT OF ATTRACTORS

Consider the product dynamics for two planar attracting homoclinic loops  $\Sigma_1, \Sigma_2 \subset \mathbb{R}^2$  (for example, the loops associated to the differential equations (7.20,7.21) described in the previous section). We investigate bifurcations that occur when we break either or both of the homoclinic connections but preserve the product structure. We will make use of the Andronov-Leontovich theorem and assume the presence of a splitting parameter (for background and more details we refer to Kuznetsov [18, §6.2], Wiggins [23] or Andronov *et al.* [3]).

FIGURE 11. Projection of  $\omega$ -limit set onto  $x_1, x_2$  plane.FIGURE 12. Projection of  $\omega$ -limit set onto  $y_1, y_2$  plane.

Specifically, for  $i = 1, 2$ , we assume that  $\xi_i$  is the splitting parameter governing the homoclinic cycle  $\Sigma_i$ , so that (see figure 13)

- (a) for  $\xi_i < 0$ , the stable manifold lies inside the unstable manifold;
- (b) for  $\xi_i = 0$ , the stable manifold coincides with the unstable manifold giving the homoclinic cycle  $\Sigma_i$ ;

(c) for  $\xi_i > 0$ , the stable manifold lies outside the unstable manifold.

Since the cycles  $\Sigma_i \subset \mathbb{R}^2$  are attracting, it follows by the Andronov-Leontovich theorem [3, 18, 23] that for sufficiently small  $\xi_i > 0$ , there exists a unique stable (hyperbolic) limit cycle  $C_i(\xi_i) \subset N_i$  such that as  $\xi_i \rightarrow 0^+$ , the limit cycle  $C_i(\xi_i)$  approaches the locus of  $\Sigma_i$  and its period  $P_i(\xi_i)$  tends to  $+\infty$ .




FIGURE 13. Phase plane as splitting parameter is varied

Keeping our earlier notation for the eigenvalues of the linearization at the equilibrium  $\mathbf{p}_i$  of  $\Sigma_i$ , we assume that

(NR)  $\frac{\mu_2}{\lambda_2} \notin \{\frac{3}{2}, 2, 3, 4\}$  and

(DI)  $\phi_t^2$  is at least  $C^7$ ,

(Conditions (NR,DI) imply that the flow  $\phi_t^2$  is  $C^3$ -linearizable at  $\mathbf{p}_2$ .)

Keeping  $\xi_2 = 0$  (so the cycle  $\Sigma_2$  persists), we break the cycle  $\Sigma_1$  by varying  $\xi_1$ . By Andronov-Leontovich theorem, there exists  $\delta_1 > 0$  such that for all  $\xi_1 \in (0, \delta)$ , there exists a unique stable limit cycle  $C_1(\xi_1) \subset N_1$ . Applying theorem 1.2 of Ashwin and Field [4], it follows that the minimal Milnor attractor for the product system  $\Phi_t^{\xi_1, 0} = (\phi_t^{1, \xi_1}, \phi_t^{2, 0})$  is the topological torus  $C_1(\xi_1) \times \Sigma_2$ .

*Remark 8.1.* Theorem 1.2 of Ashwin and Field is stated for the case when the flow  $\psi_t$  on the limit cycle is linear. That is,  $\psi_t(\theta) = \theta + \omega t$ . However, the result extends immediately to general  $C^1$ -flows on a limit cycle. Return times to a section transverse to the homoclinic loop, will be equidistributed, modulo the period. However, points of intersection with the section will generally not be equidistributed.

The torus  $C_1(\xi_1) \times \Sigma_2$  is the minimal Milnor attractor that appears along the horizontal axis of the bifurcation diagram depicted in figure 14(a). We have a similar argument when we fix the first system at  $\xi_1 = 0$  and perturb the second one.




FIGURE 14. Bifurcation diagram for the likely limit set of  $N_1 \times N_2$  for a product of homoclinic attractors.

Moreover, if we set  $\mathcal{A} = (0, \delta_1) \times (0, \delta_2)$ , then for all  $(\xi_1, \xi_2) \in \mathcal{A}$ , the likely limit set of  $N_1 \times N_2$  is the attracting normally hyperbolic two dimensional torus  $\mathbb{T}(\boldsymbol{\xi}) = C_1(\xi_1) \times C_2(\xi_2)$ . Note that  $\mathbb{T}(\boldsymbol{\xi})$  will be a minimal Milnor attractor if  $P_1(\xi_1)/P_2(\xi_1)$  is irrational — this will happen on a full measure subset of  $\mathcal{A}$ . If the ratio is rational, then the induced flow on  $\mathbb{T}(\boldsymbol{\xi})$  is a rational torus flow and so  $\mathbb{T}(\boldsymbol{\xi})$  is not even a Milnor attractor.

If the vector field associated to the flow  $\phi_t^2$  is equivariant under the group  $\mathbb{Z}_2(-I)$  generated by  $-I(x, y) = (-x, -y)$ , then the set  $\gamma(\Sigma_2) \neq \Sigma_2$  is also a homoclinic orbit associated to the origin and  $\tilde{\Sigma}_2 = \Sigma_2 \cup \gamma(\Sigma_2)$  defines a figure of eight homoclinic cycle (see section 5). Homoclinic cycles of this type appear often in the literature on the bifurcation theory of planar systems (for example, Dangelmayr & Guckenheimer [7]).

Just as above, we can break the connections for either of the homoclinic loops contained in  $\tilde{\Sigma}_2$ . Furthermore, if we analyze the first return map to a cross section in the external part of an attracting figure of eight, we can show the existence of a stable fixed point for  $\xi_2 < 0$ . That is, a periodic orbit surrounding the figure of eight bifurcates for  $\xi_2 < 0$ , as shown in figure 15 (see Guckenheimer & Holmes [11], Wiggins [23]).




FIGURE 15. Bifurcations to periodic solutions near an attracting figure of eight cycle.

Next we consider the unfolding of the product of an attracting homoclinic cycle and an attracting figure of eight (see figure 16).




FIGURE 16. Bifurcation diagram for the likely limit set of  $N_1 \times N_2$  for the product of a homoclinic attractor and an attracting figure of eight cycle.

Along the  $\xi_1$  axis (where we break the homoclinic cycle  $\Sigma_1$ ), the likely limit set consists of two two-dimensional topological tori  $C_1(\xi_1) \times \tilde{\Sigma}_2$ , which intersect in a topological circle. This will be a Milnor attractor, but not minimal. Along the positive  $\xi_2$ -axis (where we break the figure

of eight  $\tilde{\Sigma}_2$ ), the Milnor attractor has two connected components  $(\Sigma_1 \times C_{12}(\xi_2)) \cup (\Sigma_1 \times C_{22}(\xi_2))$ , each of which is a minimal Milnor attractor. Along the negative  $\xi_2$ -axis, the minimal Milnor attractor is the product of  $\Sigma_1$  and the attracting limit cycle that appears outside  $\tilde{\Sigma}_2$ . In the first quadrant of the bifurcation diagram there will be two tori which are the likely limit sets for  $N_1 \times N_2$ . These will be minimal Milnor attractors if the induced flows are irrational torus flows.

Finally, we briefly consider the product of a single homoclinic orbit and a heteroclinic attractor consisting of two equilibria and two connections. The analysis is similar to what we did for the product of two single homoclinic attractors because at least one attracting limit cycle bifurcates from the heteroclinic cycle if the interior connection is broken (see figure 17). Given the heteroclinic network shown in figure 17, a characteristic situation is that there is a unique unstable equilibrium enclosed by each cycle (when  $\xi = 0$ ). When we break the interior heteroclinic connection from  $A$  to  $B$ , an attracting limit cycle is created. We show a typical scenario in figure 17.




FIGURE 17. An unfolding of a heteroclinic cycle.

## 9. DISCUSSION AND CONCLUSION

Motivated by the partial results in Ashwin and Field [4] and supported by numerical simulations, we have proved that the likely limit set of the product of two planar heteroclinic attractors is the one-dimensional heteroclinic network which covers the attracting networks in the factors. The result implies that generically two independent trajectories around the heteroclinic connections are forward asynchronous. Following the analysis presented here, it is not difficult to prove that the likely limit set of the product of a finite number of heteroclinic attractors is also the 1-dimensional heteroclinic network which covers the attracting networks in the factors.

For the proof of our main result we needed to assume that the flow was at least  $C^2$ . The situation when the flow is only  $C^1$  is far from clear, especially in the resonant case when  $\rho_1 = \rho_2$ . Since theorem 2.6 holds for  $C^1$ -flows, a counterexample in the  $C^1$ -case would give the likely limit set as the product of heteroclinic attractors and would likely depend on large fluctuations in the derivative of the connection maps. Similar issues of differentiability arise in the case of the product of a homoclinic attractor and a limit cycle (the result in Ashin and Field [4] required the homoclinic flow to be  $C^7$ ) and it is not clear whether or not the product of a homoclinic attractor and a limit cycle is a minimal Milnor attractor if the flow (for the homoclinic factor) is less regular, for example  $C^2$ .

In the context of game theory and the replicator equation, Sato *et al.* [21] studied numerical examples of heteroclinic networks in the product of simplices. In the case of the Rock-Scissors-Paper game, the dynamics is dependent on the payoff matrices between the players. Under some conditions, numerical evidence was found for the existence of complicated behaviour near the product network. There remain questions about the heteroclinic network which appears in the product dynamics. One candidate heteroclinic network for the product is the product of the heteroclinic networks for each simplex. However, only some cycles might be observable and these may depend on the interaction between the players. This is one reason why we believe our result may have interesting applications outside of equivariant dynamics and network dynamics [1, §5] and why it would be worthwhile generalizing to products of two heteroclinic cycles in higher dimensions. In dimensions greater than 2, issues such as the orientability of the homoclinic connection and the existence of degenerate cases of homoclinic cycles, such as orbit flip and inclination flip, complicate the study. There is also the question of considering the product of a homoclinic cycle with a *butterfly* or *bellows*. We refer to the recent survey by Homburg and Sandstede [14] for more details on these types of cycle as well as issues connected to bifurcation theory and the breakdown of hyperbolicity at the saddle point.

The outstanding question is undoubtedly to obtain quantitative results about the effects of perturbations breaking the product structure to, for example, a skew product structure but keeping the heteroclinic cycles. Does the loss of the product structure lead to phenomena related to essential asymptotic stability or do the likely limit set results persist for small enough perturbations? Put another way, is it possible to find cycle preserving perturbations that make a pre-specified subnetwork essentially asymptotically stable?

## REFERENCES

- [1] M Aguiar, P Ashwin, Dias and M Field. ‘Dynamics of coupled cell networks: synchrony, heteroclinic cycles and inflation’, *Journal of Nonlinear Science*, to appear.
- [2] M A D Aguiar, I S Labouriau and A A P Rodrigues. ‘Switching near a network of rotating nodes’, *Dynamical Systems: an International Journal* **25**(1) (2010), 75–95.
- [3] A A Andronov, E A Leontovich, I I Gordon, A G Maier. *Theory of Bifurcations of Dynamical Systems on a Plane* (Israel Program of Scientific Translations, Jerusalem, 1973).
- [4] P Ashwin and M Field. ‘Product dynamics for homoclinic attractors’, *Proceedings of Royal Society, Series A* **461** (2005), 155–177.
- [5] P Ashwin and P Chossat, ‘Attractors for Robust Heteroclinic Cycles with Continua of Connections’, *J. Nonlinear Sci.* **8** (1998), 103–129.
- [6] W Brannath. ‘Heteroclinic networks on the tetrahedron’, *Nonlinearity* **7** (1994), 1367–1384.
- [7] G Dangelmayr and J. Guckenheimer, ‘On a four parameter family of planar vector fields’, *Arch. Rational Mech. Anal.* **97** (1987), 321–352.
- [8] B Deng. ‘The Shilnikov problem, exponential expansion, strong  $\lambda$ -lemma,  $C^1$ -linearization, and homoclinic bifurcations’, *Journal of Differential Equations*, **79** (1989), 189–231.
- [9] B Ermentrout. *Simulating, Analyzing, and Animating Dynamical Systems: A Guide to XPPAUT for Researchers and Students* (SIAM, Philadelphia, USA 2003).
- [10] M Field. *Dynamics and Symmetry* (Imperial College Press Advanced Texts in Mathematics – Vol. 3, 2007.)
- [11] J Guckenheimer and P Holmes, ‘Structurally stable heteroclinic cycles’, *Math. Proc. Camb. Phil. Soc.* **103** (1988), 189–192.
- [12] P Hartman. ‘On local homeomorphisms of Euclidean spaces’, *Bol. Soc. Mat. Mexicana* **2**(5) (1960), 220–241.
- [13] A J Homburg and J Knobloch. ‘Switching homoclinic networks’, *Dynamical Systems* **23** (2010), 351–358.
- [14] A J Homburg and B Sandstede. ‘Homoclinic and Heteroclinic Bifurcations in Vector Fields’ (Chapter 8 in *Handbook of Dynamical Systems*, vol. 3, Elsevier, 2010).
- [15] J F C Kingman and S J Taylor. *Introduction to Measure and Probability*, (Cambridge University Press, 1966).
- [16] V Kirk and M Silber. ‘A Competition Between Heteroclinic Cycles’, *Nonlinearity* **7** (1994), 1605–1621.
- [17] M Krupa and I Melbourne. ‘Asymptotic Stability of Heteroclinic Cycles in Systems with Symmetry, II,’ *Proc. Roy. Soc. Edinburgh* **134A** (2004), 1177–1197.
- [18] Yuri A Kuznetsov. *Elements of Applied Bifurcation Theory* (Applied Mathematical Sciences **112**, Springer-Verlag, 1995).
- [19] I Melbourne. ‘An example of a non-asymptotically stable attractor’, *Nonlinearity* **4** (1991), 835–844.
- [20] J Milnor. ‘On the concept of attractor’, *Commun. Math. Phys.* **99** (1985), 177–195.

- [21] Y Sato, E Akiyama and J P Crutchfield, ‘Stability and Diversity in Collective Adaptation’, *Physica D* **210** (2005), 21–57.
- [22] L S Shilnikov. *Methods of Qualitative Theory in Nonlinear Dynamics, Part II* (World Scientific Publ. Co. Ltd., Singapore, 2001).
- [23] S Wiggins. *Global Bifurcations and Chaos: Analytical Methods* (Springer-Verlag, 1988).
- [24] S Wiggins. *Introduction to Applied Nonlinear Dynamical Systems and Chaos* (Springer-Verlag, 2003).

NIKITA AGARWAL, DEPARTMENT OF MATHEMATICS, UNIVERSITY OF HOUSTON, HOUSTON, TX 77204-3008, USA

*E-mail address:* `nagarwal@math.uh.edu`

ALEXANDRE RODRIGUES, CENTRO DE MATEMÁTICA DA UNIVERSIDADE DO PORTO AND FACULDADE DE CIÊNCIAS DA UNIVERSIDADE DO PORTO, RUA DO CAMPO ALEGRE 687, 4169-007 PORTO, PORTUGAL

*E-mail address:* `alexandre.rodrigues@fc.up.pt`

MICHAEL FIELD, DEPARTMENT OF MATHEMATICS, UNIVERSITY OF HOUSTON, HOUSTON, TX 77204-3008, USA

*E-mail address:* `mf@uh.edu`